Index to Volume 82 of the Bryologist[1,2]

Absconditella sphagnorum, 316
Acarospora chlorophana, 22; fuscata, 22, 24, 125, 131; heppii, 125, 131; schleicheri, 22, 131; smaragdula, 22; strigata, 22, 24; sulphurata var. rubescens, 504
Acaulon muticum, 519, 542, 554, var. rufescens, 542; schimperianum, 542; triquetrum, 542
Acrophyllum javense, 498
Acroschisma wilsonii, 514
Acidodontium megalocarpum, 228; rhamphostegium, 228
Acrocladium auriculatum, 514; politum, 246
Acrocryphaea gardneri, 234, 435; leiboldii, 434
Acromastigum longirete, 311
Acroporium pungens, 242, 481
Actinodontium sprucei, 237
ACTON, D. W. *See* J. M. GLIME & D. W. ACTON, 382
Adelothecium bogotense, 237, 439, 443
Additional new and noteworthy records for Hawaiian mosses—5, 79
Agonimia octospora, 315; tristicula, 500
air pollution, 20+, 54+, 280+
Alaska, Lichens of Anaktuvuk Pass, 393; Molendoa hornschuchiana new to, 487
Alberta, Bryophytes new to, 409
Alectoria irvingii, 396+; lanea, 403; nigricans, 402; ochroleuca, 108, 397+
ALLEN, B. H. Additional distribution records of mosses in Pennsylvania, 289
ALLEN, B. H. Bryoxiphium norvegicum in Colorado, 490
Aloina aloides, 542, 554, var. ambigua, 261, 542; bifrons, new to South America, 493; brevirostris, 542; hamulus, 629; pilifera, 261; rigida, 72, 261, 542
Aloinella catenula, 629
ALPERT, P. Desiccation of desert mosses following a summer rainstorm, 65
Alpine mosses of Mexico, Notes on, 629
Alsia californica, 261
Amblystegiella confervoides, 290; subtilis, 622
Amblystegium fluviatile, 484; juratzkanum, 261, 611, 622; riparium, 525+, 556, forma longifolium, 484, Temperature related growth patterns in, 525; serpens, 261, 274, 492, 514, 611; tenax, 484; varium, 261, 274, 484, 492, 611, var. ovatum, 611, var. parvulum, 609, 610
Amblytropis gemmacea, 237; hispidula, 237; ovata, 237
amino acids in liverworts, 287+
Amphidium californicum, 261; cyathicarpum, 232, 629; lapponicum, 79, 519
Amphoritheca jamesonii, 246; lindigii, 228, 246; lutescens, 228, 246; pseudomarginata, 228, 246; undulata, 228, 246
Anacalypta cucullata, 227, 246; subcaespitosa, 226, 246
Anacamptodon cubensis var. latifolius, 494; mexicanus, 494; splachnoides, 492
Anacolia intertexta, 434; laevisphaera, 230, 246, 247, 249; menziesii, 261; subsessilis, 230, 246; webbii, 515
Anaptychia diademata, 473, forma brachyloba, 473; neoleucomelaena, 564; obscurata, 473
Anastrophyllum assimile, 163+, 413, 415, 457, 467; ellipticum, 634; minutum, 166
Ancistrodes genuflexa, 518
ANDERSON, L. E. *See* H. A. CRUM & L. E. ANDERSON, 469
Andreaea aristata, 220, 246; brevipes, 220; nitida, 220; obovata, 168; rupestris, 80, 220, 496, 514; subenervis, 220; subulata, 220
Andreaeobryum macrosporum, 374, 409, 416
ANDRUS, R. E. Sphagnum bartlettianum in the Southeastern United States, 198
ANDRUS, R. E. & D. H. VITT. Sphagnotheca boreali-americana: Fascicle III, 295
Aneura pinguis, 582
Anisothecium convolutum, 221; hookeri, 221, 246; jamesonii, 221, 246; vaginatum, 221, 246
Anoectangium aestivum, 375, 544, 554; incurvans, 434; obtusifolium, 261, 484; sendtnerianum, 484; stracheyanum, 544; thompsonii, 544
Anomobryum clavicaule, 228; filiforme, 228, 246, 247, 413, 438, 489; julaceum, 515; plicatum, 228; prostratum, 228, 246; semiovatum, 228
Anomodon angustatus, 246; attenuatus, 274, 438, 442, 484, 611; dentatus, 496; minor, 274, 484, 611; rostratus, 74, 142, 274, 438, 442, 484, 492, 611; scaberrimus, 16; viticulosus, 71, 484
Anomomarsupella cephaloizielloides, 163
antagonism of cytokinin induced callus in Py-

---

[1] Compiled by William C. Steere, New York Botanical Garden, Bronx, New York 10458.

[2] New scientific names are printed in boldface types; illustrations are indicated by an asterisk; taxa which are discussed or mentioned several times in a paper are indicated with a plus sign.

laisiella selwynii by nucleosides and cyclic nucleotides, 47
Anthelia juratzkana, 163+
Anthoceros adscendens, 298+; erectus, 145, 146, 151
Anthracothecium goniostomum, 473; inclinatum, 503
antibiotic activity of bryophytes, 141
antibiotic substances, 141+
Antitrichia californica, 261; curtipendula, 261
Anzia jamesii, 316
Aongstroemia acerosa, 221, 246; bogotensis, 223, 246; callosa, 223, 246; consimilis, 223, 246; convoluta, 221, 246; crassinervis, 223, 246; curviseta, 221, 246; ditissima, 223, 246; julacea, 221; longipes, 516; maculata, 221; muralis, 224, 246; pusilla, 224, 246; strumulosa, 224, 246, var. minor, 224; vaginata, 246
Apalodium pellucens, 246
Aphanorrhegma serratum, 611
Aptychella colombica, 242; proligera, 242
Aptychopsis pycnodonta, 242
aquatic bryophytes, zinc localization in, 594
archegonium and embryo in Lophocolea heterophylla, 576
ARCHER, A. W. & W. A. WEBER. Baeomyces frenchianus, an Australian endemic and its chemotaxonomy, 481
Archidium acauloides, 498; phascoides, 514; tenerrimum, 492
Argentina, Scouleria patagonica in, 99+
Arizona, desiccation of desert mosses in, 165
Arkansas, Bryophytes of the Ozark National Forest, 483
Arthonia atlantica var. positiva, 315; caesia, 125, 131; swackhii var. macrospora, 315
Arthothelium illicinum var. dictyosporum, 501, var. reagens, 501; lirellans, 501
Arthrorhaphis citrinella, 402
Asahinea chrysantha, 397+; scholanderi, 396+
Aschisma carniolicum, 546
Aspidothelium trichothelioides, 110
Aspilicia aquatica, 395+; caesiocinerea, 22, 402; calcarea, 22; chadefaudiana, 319; cinerea, 22, 402; nikrapensis, 402; plicigera, 395+; sublapponica, 395+; supertegens, 402
Aspiromitus cavernosus, 118; punctatus, 118
assimilation in Fontinalis duriaei, 382
Asterella angusta, 148, 151; elegans subsp. echinella, 299; gracilis, 163, linenbergiana, 163; sanguinea, 141, 145, 148, 151; tenella, 299, 483
Asteriscium decolorans, 227, 246
Asterothyrium argenteum, 89; leucophthalmum, 88+, 505
Astomum crispum, 545, var. sterile, 545; exertum, 545; lindigii, 225; ludovicianum, 492, 540, 545; mittenii, 545; muhlenbergianum, 540, 545, 609, 611
Aswathia melanotricha, 107
Athalamia hyalina, 163
Atractylocarpus costaricensis, 434+; longisetus, 221, 222, 248; spiripes, 221; strictulus, 495
atranorin, 195, 292+, 482
Atrichopsis magellanica, 519
Atrichum altecristatum, 609, 610; androgynum, 244, 246; angustatum, 29, 30, 32, 33, 274, 438, 484, 609, 611; bogotense, 244, 246; oerstedianum, 244, 434, 609, 610; polycarpum, 245, 434; rhystophyllum, 438; tenellum, 609, 610; undulatum, 93+, 142, 274
Aulacomnium acuminatum, 410; androgynum, 74, 261; heterostichum, 484, 609, 611; palustre, 74, 83, 261, 328, 438, 484, 492, 496, 611; turgidum, 496
Aluaxina microphana, 88, 91, 131; quadrangula, 125, 131; unispora, 110
Austinia brotheri, 494+; tenuinervis var. micholitzii, 494

Bacidia alutacea, 473; bagliettoana, 402; chlorococca, 25, 125, 131; coprodes, 396+; fusca, 402; inundata, 125, 131; lambinonii, 110; lugubris, 402; melaena, 402; michaeliana, 110; microcarpa, 402; naegelii, 125, 131; obscurata, 402; schweinitzii, 131; sphaeroides, 402; subincompta, 396+; vezdae, 315
Baeomyces frenchianus, an Australian endemic and its chemotaxonomy, 481+; trachypus, 481
Baeomycesic acid, 154+
BAKER, K. K., J. A. MALACHOWSKI & G. R. HOOPER. Ultrastructural localization of polysaccharides in Usnea cavernosa, 533
BANERJEE, R. D. & S. P. SEN. Antibiotic activity of bryophytes, 141
barbatic acid, 154+
Barbella pendula, 440; tenuissima, 234
Barbilophozia barbata, 298+; quadriloba, 118
Barbula acuta, 72; affinis, 227, 246; agraria, 541; andreaeoides, 413; apiculata, 226, 246; appressifolia, 226; aurea, 499; bicolor, 543; bogotensis, 227, 246; brachyphylla, 261; campylocarpa, 226; cancellata, 609, 611; convoluta, 261, 375, 543, 609, 610; coreense, 543; crocea, 543; decolorans, 227, 246; enderesii, 543; fallax, 275, 289; fragilis, 227, 246; glaucescens, 226; goudotii, 227, 246; horridinervis, 543, 549, 551; inaequalifolia 226, 259, 619; indica, 541; laevigata, 226; lindigii, 246; longirostris, 226; nigrescens, 413; novo-granatensis, 226, var. gracilior, 226, var. teretiuscula, 226; orizabensis, 434; pringlei, 434; rectifolia, 226; replicata, 226, 246, 255; revoluta, 543, var. hornschuchiana, 543; rubiginosa, 261, 499; stenocarpa, 226; subcaespitosa, 226; trian-

ae, 226, 246; unguiculata, 261, 275, 484, 519, 543, 611; vinealis, 261
BARCLAY-ESTRUP, P. & R. J. K. RINNE. Trace element accumulation in a feather moss and in soil near a Kraft paper mill in Ontario, 599
Bartramia angustifolia, 230; bogotensis, 231, 246; chrysea, 231, 246; curvata, 231, 246; defoliata, 230; dilatata, 230; elegantula, 246; flavicans, 231, 246, 247; fontanella, 231, 246; fragilifolia, 231; gracilenta, 232, 246; halleriana, 515; humilis, 231, 246; jamesonii, 231, 246; karsteniana, 231, 246; lindigii, 246; longifolia, 231; lutescens, 231, 246; macrotheca, 231, 246; mathewsii, 231; minuta, 232, 246; patens, 99; pinnata, 232, 246; polytrichoides, 231; pomiformis, 275, 484, 611; potosica, 231; squarrosa, 231, 246; striatula, 232, 246; stricta, 261; strumosa, 231; subarcuata, 247; subpatens, 231, 247; subsessilis, 230; tenella, 232, 247; thelioides, 231; tomentosa, 247; trianae, 231, 247; versifolia, 232, 247
Bartramidula lindigii, 231; wilsonii, 515
Bazzania aterrima, 116; indigenarum, 116; minutiserra, 116; pompeana, 117; trilobata, 274; vittata, 634
Bestia longipes, 261; vancouveriensis, 261
Bilimbia microphana, 88
Bizotia densifolia, 223, 247
Black Hills of South Dakota and Wyoming, addition to the mosses of, 72
Blepharostoma trichophyllum, 163+
Blindia acuta, 221, 520, 583; campylopodioides, 583, 584; inundata, 247; japonica, 583; magellanica, 221, var. inundata, 247; sonsoniae, 221
Bombyliospora vulpina, 125, 131
BOWERS, F. D. *See* COLE, M., F. D. BOWERS & S. FRECKMANN, 273
Brachydontium flexisetum, 221
Brachymenium columbicum, 247; consimile, 228, 434; daguense, 228; exile, 438; indicum, 79, 81; morasicum, 228; nepalense, 144; papillosum, 247; speciosum, 228, 253; spirifolium, 434, 441; systylium, 435, 441, 443
Brachymitrion jamesonii, 228; moritzianum, 228, 255
Brachyodus flexisetus, 221, 247
Brachysteleum lindigii, 232, 247
Brachythecium acuminatum, 275, 484, 492, 609, 611; acutum, 609, 611; albicans, 261; asperulum, 240; bolanderi, 262; calcareum, 484; chrysocladon, 240; collinum, 72, 262; conostomum, 241; flexiventrosum, 241; frigidum, 262; groenlandicum, 413; kamouense, 146; laevifolium, 241; mildeanum, 413; nelsonii, 412; occidentale, 241; oxycladon, 275, 484, 609, 611; plumosum, 74, 241, 275, 438; prasiophyllum, 241; procumbens, 141, 145, 147, 150, 151; pseudorutabulum, 241; reflexum, 275, 609, 610; rivulare, 87, 275, 611; rutabulum, 87, 241; salebrosum, 275, 611; serrulatum, 440; stereopoma, 241; velutinum, 262, var. venustum, 262
BRASSARD, G. R. (review) T. Koponen, P. Isoviita & T. Lammes, The bryophytes of Finland: an annotated checklist, 121
BRATT, G. C. *See* RUNDEL, P. W., G. C. BRATT & O. L. LANGE, 171
Breutelia allionii, 231; arcuata, 515; brittoniae, 231; chrysea, 231, 247; falcatula, 231; inclinata, 231; integrifolia, 231; jamaicensis, 435; karsteniana, 231; lorentzii, 231; rhytidioides, 231; sphagneticola, 231, 247; squarrosa, 231; subarcuata, 231; subplicata, 523; tomentosa, 231, 247; trianae, 231
British Columbia, new and interesting hepatics from, 162+
British Honduras, some mosses from, 478
Brothera leana, 484
Brotherella recurvans, 275, 491; tenuirostris, 275
Bruchia lindigiana, 221; subenervis, 221
Bryhnia graminicolor, 72, 275, 484, 611; novaeangliae, 275
Bryoandersonia illecebra, 484, 492, 611
Bryobrittonia longipes, 410, 450+; pellucida, 416, 456+
Bryobrothera crenulata, the systematic position of, 305+, 307*
Bryoerythrophyllum alpigenum, 544; angustulum, 247; arcuatum, 226; brachystegium, 544; campylocarpum, 226, 255; ferruginascens, 413; gymnostomum, 544; jamesonii, 226, 487; recurvirostre, 74, 262, 519; recurvirostrum, 275, 461, 544, 609, 611; rubrum, 544, var. minus, 544; yichunense, 496
Bryomanginia radiculosa, 498; saint-pierrei, 631
Bryoria nitidula, 403
Bryosedgwickia densa, 439
Bryoxiphium norvegicum, 274+, in Colorado, 490, var. mexicanum, 629
Bryum albicans, 229, 247; algovicum. 72, 611; antarcticum, 392, 532; angustirete, 262; apiculatum, 80; argenteum, 228, 247, 262, 392, 438, 442, 484, 515, 532, 609, 611; aubertii, 619; beyrichianum, 247; bicolor, 262, 484, 498; billardieri, 229, 247, 440; bimum, 229, 247; bulbillosum, 229, 247; caespiticium, 262, 275, 611; canariense, 262; candicans, 229; capillare, 67+, 262, 275, 411, 416, 438, 484; clavicaule, 247; coronatum, 229, 480; corrugatum, 228, 247; creberrimum, 73, 262, 611; cruegeri, 229, 480, 560; dentatum, 434, 441; dichotomum, 247; dunense, 498; ellipsifolium, 619; erythrocar-

pum, 79, 82; erythroneuron, 229; flaccidum, 413; flavituber, 79, 82; fragilicuspis, 229; gemmiparum, 73, 484; genucaule, 619; goudotii, 229, 247; grandifolium, 247; intermedium, 413; julaceum, 228, 247; klinggraeffii, 79, 497; laevigatum, 229; leptodontium, 230, 247; leptochaeton, 229; limbatum, 435; lindigianum, 229; lisae var. cuspidatum, 275, 289; lonchocaulon, 262; mayorii, 229; megalocarpum, 247; miniatum, 262; mniopsis, 230, 247; pallescens, 262; papillosum, 247; paucifolium, 230, 247; perconcavifolium, 434, 441; procerum, 229, 454; proliferum, 229; prostratum, 228, 247; pseudotriquetrum, 229, 247, 262, 484, 609, 611; pycnopyxis, 247; radiculosum, 440; remotifolium, 229; rhamphostegium, 247; rubens, 495; salaminae, 230, 247; semimarginatum, 229; semiovatum, 247; socorrense, 229; sordidum, 229; spectabile, 229, 247; stenotrichum, 262; subcurvatum, 230, 247; terebellum, 229; truncorum, 229, 247, 258; turbinatum, 262, var. minus, 229, 247; uliginosum, 73; weigelii, 82, 83
Bucklandia bartrami, 523
Buellia alboatra, 22; amphidexia, 132; buellioides, 500; callispora, 125, 132, 318; curatellae, 132; elizae, 125, 132; leucomela, 132; malmei, 396*; moriopsis, 403; novomexicana, 22; papillata, 403; punctata, 22, 403; var. polyspora, 125, 132; pulverea, 315; retrovertens, 22; scabrosa, 403; spuria, 132; stibmaea, 132; zahlbruckneri, 22
Bulbothrix goebelii, 135; laevigatula, 135
BURTON, M. A. S. & P. J. PETERSON. Studies on zinc localization in aquatic bryophytes, 594
Buxbaumia aphylla, 29, 36
Byssoloma leucoblepharum, 88, 91, 125, 132; rotuliforme, 88; subdiscordans, 88, 91; tricholomum, 88; vezdanum, 110

Calenia africana, 110; conspersa, 110
California, preliminary list of southern California mosses, 260; photosynthetic response to lichens to short-term ozone fumigations, 280
Callicladium haldanianum, 611
Callicosta andersonii, 494; armata, 494; asperifolia, 237, 494; bipinnata, 237; corrugata, 494; cristata, 494; debilis, 494; divaricata, 494; fendleri, 237, 494; husnotii, 494; longicaulis, 494; lophophylla, 494; luciae, 494; procera, 494; ramosissima, 237, 253, 494; rugifolia, 494
Callicostella acutifolia, 237; colombica, 237; depressa, the chromosome number of, 103; pallida, 237; plicatula, 237
Calliergidium bakeri, 514
Calliergon cordifolium, 514; cuspidatum, 240; sarmentosum, 240
Calliergonella cuspidata, 247
callus in Pylaisiella selwynii, 47+
Calobryum gibbsiae, 582
Calomnium dixonii, 306
Caloplaca amabilis, 22; arenaria, 22; arizonica, 22; atroalba, 22; aurantia, 501; castellana, 318; cerina, 22, 403; cinnamomea, 403; decipiens, 22; diphyodes, 22; epithallina, 22; ferruginea, 22, 110; flavovirescens, 125, 132; **floridana**, 125, 132; fraudans, 403; holocarpa, 22, 403; intermedia, 125, 132; isidiigera, 506; jungermanniae, 493; microphyllina, 22; modesta, 22; pinicola, 22, 493; stillicidiorum, 403; tetraspora, 403; tiroliensis, 403; trachyphylla, 22, 24, 25, 27; xanthaspis, 473
caloplacin, 302+
Calymperes bartramii, 225; donnellii, 225, 479, 559, 560; erosum, 225, 559, 560; guildingii, 619; lindigii, 227, 247; lonchophyllum, 225, 479, 560; nicaraguense, 560; richardii, 225, 479, 561; uleanum, 225, 561
Calypogeja fissa subsp. neogaea, 483; meylanii, 113; muelleriana, 118, fo. paludosa, 118, fo. sphagnicola, 118, fo. tundrae, 118; neesiana var. meylanii, 118; sphagnicola, 118; suecica, 118; trichomanis, 610
Calyptothecium duplicatum, 236, 435; moritzii, 236
Calyptrochaeta enervis, 498; flaccida, 498; parviretis, 498; rotundifolia, 498
Camptochaete arbuscula, 518
Camptodontium brotheri, 516
Camptothecium pseudo-lutescens, 241
Campylium chrysophyllum, 275, 438, 485, 611; hispidulum, 240, 275, 611; protensum, 495; radicale, 413
Campylopodiella ditrichoides, 585, 587
Campylopodium curvisetum, 221; euphorocladum, 516; pusillum, 435
Campylopus albidovirens, 618; altissimus, 247; annotinus, 221; apollinairei, 221; arctocarpus, 221; areodictyon, 221; argyrocaulon, 221, 223; artocarpus, 435, 442; atrovirens, 585; benedictii, 221, 618, 623; bogotensis, 221; breweri, 221; capitulatus, 618; cavifolius, 222; chionophilus, 222, 618; chrismarii, 222, 247; chrysodictyon, 222; concolor, 222, 258; cuatrecasii, 222; cucullatifolius, 222; densicoma, 222, 611; filicaulis, 222; filicuspis, 222, 618; filifolius, 222, 247, 248; flexuosus, 222, 479, 516, 585; fragilis, 222, 585; friabilis, 222, 247; fulvus, 618; galapagensis, 495; goughii, 585, 587; gracilis, 144, 585; harpophyllus, 222; hawaiicus, 80; heterophyllus, 222; heterostachys, 434, 441; huallagensis, 222; incacor-

ralis, 618; incertus, 222, 618; incurvatus, 495; introflexus, 223, 257, 448, 485, 497; jamesonii, 222, 247, 248, 258; jugorum, 222; kingii, 222; laetus, 146, 151, 152, 585; lamellinervis, 222, 247; lamprodictyon, 222, 434; latinervis, 585; leucognodes, 222; longisubulatus, 619, 623; mexiae, 222; oblongus, 222; occultus, 222; paramoensis, 222; pauper, 222, 247, 347, 619, var. minor, 223; penicillatus, 222, 247; pilifer, 223, 247, 257, 438; pittieri, 221, 223, 247, 253; polytrichoides, 223, 247, 257, 448; porphyreodictyon, 222, 247; praemorsus, 80; propinquus, 222, 223, 247; reflexisetus, 223; revolvens, 223; richardii, 144, 223, 253, 255, 585; rosulatus, 223; savannarum, 223, 434; schimperi, 409, 415, 495; **sharpii**, 623, 624*, 625; strictulus, 223; subconcolor, 223, 619, 629; subfimbriatus, 619; subfulvus, 619; subulatus, 415, 495; surinamensis, 223; tequendamensis, 223; thysanomitrioides, 223, 619; trichophorus, 223; tunariensis, 223; walkeri, 585; ynesiae, 222; yungarum, 223

Canada, bryophytes new to, 409

Candelaria concolor, 473, var. vainioana, 473

Candelariella athallina, 396+; aurella, 23, 403; deflexa, 23; efflorescens, 316; lutella, 306+; rosulans, 23, 24; terrigena, 493; vitellina, 493

Carbon dioxide fixation rate, 171+

Caribou grazing, impact on lichens of Anaktuvuk Pass, Alaska, 393

Catagoniopsis berteroana, 515

Catagonium politum, 236, 250, 519, var. phyllogonium, 236

Catharinea abriaquiae, 245, 247; aristata, 245, 247; ciliata, 247; erosa, 245, 247; polycarpa, 247; trianae, 245, 247; trichodon, 247

Catillaria bouteillei, 88, 125, 132, 505; kansuensis, 23

Catolechia wahlenbergii, 502

Cavicularia densa, 117

Cephalozia bicuspidata, 274; heteroica, 310; kashyapii, 637; laxifolia, 637; lunulifolia, 274, 299; pleniceps, 610; zoopsioides, 115

Cephaloziella elastica, 116; hampeana, 163; rubella, 299; uncinata, 633; violacea, 118

Ceratodon dimorphus, 476; novo-granatensis, 220; purpureus, 50, 220, 262, 275, 461; propaguliferous, 474+, 516, 609, 611, subsp. dimorphus, 477, var. dimorphus, 475, 477, forma dimorphus, 474, 477; stenocarpus, 220; venezuelensis, 220

Cetraria andrejevii, 493; cucullata, 318, 398+; delisei, 110, 401+; ericetorum, 493; hepatizon, 493; islandica, 397+; laevigata, 493; nigricans, 396, 493; nivalis, 318, 319, 398+, 493; pinastri, 493; sepincola, 493; tilesii, 493

Cetrelia cetrarioides, 501; monachorum, 501

Chaenotheca laevigata, 506; xyloxena, 506

Chaenothecopsis fennica, 320; parasitaster, 502

Chaetomium mollicellum, 319

Chamberlainia stereopoma, 435

Chandonanthus setiformis, 168

Cheilolejeunea acutangula, 634; inflexa, 634

Cheilothela boliviana, 220

chemical analyses of Lecidea, 189

chemosyndromes of Xanthoparmelia, 154

chemotaxonomy of Baeomyces frenchianus, 481+

Chile, Aloina bifrons new to, 493; mosses of, 99+

Chiloscyphus pallescens, 581, var. fragilis, 483

Chionodecton perplexum, 125, 133; montagnaei, 133; sanguineum, 133

chlorodepsidones, 302

chlorophyll content of liverworts, 286+

Chorisodontium aciphyllum, 361, 365; speciosum, 223; wallisii, 223

chromosome number of Callicostella depressa, 103

CHURCHILL, S. P. Mosses of the Great Plains III. Additions to Nebraska and the Black Hills of South Dakota and Wyoming, 72

Cinclidotus aquaticus, 546; fontinaloides, 546, 554; riparius, 546

Cladina macronesica, 108; mediterranea, 108; stellaris, 564

Cladodium rhamphostegium, 228, 247; socorrense, 247

Cladomniopsis crenato-obtusa, 520

Cladonia alaskana, 396; alpestris, 110, 398, 408; alpina, 320, 493; amaurocraea, 318, 493; apodocarpa, 133; arbuscula, 284, 393+, 504, 564, 575; bacilliformis, 493; bellidiflora, 493; caespiticia, 316; cariosa, 493; caroliniana, 133; chlorophaea, 133, 493; clavulifera, 133; coccifera, 493; coniocraea, 23; cornuta, 493; crispata, 396; deformis, 493; degenerans, 493; emocyna, 179, 493; fimbriata, 493; gonecha, 493; gracilis, 396+, impexa, 493; invisa, 316; lepidota, 493; luteoalba, 407, 493; macroceras, 314; macrophylla, 493; mateocyatha, 133; maxima, 314; metacorallifera, 493; mitis, 71, 180, 393+, 504, 574; norrlinii, 396+; phyllophora, 493; pleurota, 493; pocillum, 493; portentosa forma subimpexa, 314, subsp. pacifica, 314, forma decolorans, 314; pseudostellata, 396+; pyxidata, 23, 404; rangiferina, 71, 109, 180, 393+; rangiformis, 105, 283, 314, 564, 574, var. gracillima, 314; squamosula, 482; stellaris, 284+, 393+, 503, 575; stricta var. uliginosa, 314; subulata, 404; subtenuis, 180; tenuis, 396, 404; thomsonii, 105, 404;

uncialis, 71, 180, 404; verruculosa, 105; verticillata, 404; wainii, 105
Clapodium subpiliferum, 168; whippleanum, 262
Clasmatodon parvulus, 485, 492
Climacium americanum, 275, 485, 611, 622; dendroides, 74, 275
Cnestrum alpestre, 496; schistii, 496
coal-fired electricity generating facilities on N_2-fixing lichens, impact of emissions from, 54
Coccocarpia pellita, 396+
cold injury to lichens in southern Ohio, 620
COLE, M., F. D. BOWERS & S. FRECKMANN. Bryophytes of the Kickapoo River Valley, southwestern Wisconsin, 273
Coleochaetium scaberrimum, 16; standleyi, 16
Collema ceraniscum, 404; coccophorum, 23; furfuraceum, 23; glaucophthalmum, 473; glebulentum, 404; multipartitum, 396+; nigrescens, 564+; polycarpon, 404; tenax, 23, 54+, 316, 505; tunaeforme, 23, 404; undulatum, 23, 404, var. granulosum, 23
Cololejeunea aoshimensis, 118; biddlecomiae, 483, 610; cristata, 604; cuneata, 604; decemplicata, 606; **diplasiolejeunoides**, 602, 603*; **disciflora**, 602, 604*; distallo-papillata, 608; falcata, 608, var. **madecassa**, 602, 605*, 606; minutissima, 605; occidentalis, 313; rossettiana, 119; **sambiroana**, 602, 607*, 608; Some new species of Malagasy, 602
Colorado, Bryoxiphium norvegicum in, 490
Colura calyptrifolia, 312; tenuicornis, 312
Coniocybe coniophaea, 506; peronella, 320
Conocephalum conicum, 74, 96, 97, 98, 112, 141, 142, 147, 150, 151, 274, 483, 610, 634; supradecompositum, linear spore tetrads in, 96, 97*, 98*, 634
Conomitrium intromarginatum, 220, 247; lindigii, 220
Conostomum arcticum, 515; jamesonii, 231, 246; pentastichum, 231
contribution to the lichen flora of Uruguay XI. Lichens from the Arazatí, San José Department, 472
Cornicularia aculeata, 404; divergens, 398+
Costesia spongiosa, 517
CRANDALL-STOTLER, B. *See* R. E. STOTLER & B. CRANDALL-STOTLER, 112, 310, 632
Crater Lake National Park, Oregon; new mosses from, 82
Cratoneuron filicinum, 145, 146, 240, 262, 485, 514
CROSBY, M. R. Recent literature on mosses—113, 494
Crossidium aberrans, 542; desertorum, 262; squamiferum, 519, 542, var. pottioideum, 262, 542
Crossocalyx koriakensis, 118
Crossogyna autumnalis, 118; elongella, 118; flexicaulis, 118; kirkii, 118; nipponica, 118; orbicularis, 118; purpurascens, 118; robusta, 118; tasmanica, 118; undulifolia, 118
Crossomitrium oerstedianum, 237; patrisiae, 237; wallisii, 237
CRUM, H. A. & L. E. ANDERSON. Tortula ammonsiana, a new species from West Virginia, 469
Crummia latifolia, 262
Cryphaea cuspidata, 234, 247; gardneri, 247; glomerata, 485; lindigiana, 234, 247; lindigii, 247; novo-ranatensis, 234; patens, 234, 434; pilifera, 234; tenuissima, 248
Cryphaeophyllum molle, 518, 522
Cryptopodium hookeri, 248
Cryptothallus mirabilis, 311
Ctenidium malacodes, 243, 435; molluscum, 485
CULBERSON, C. F. & H. HERTEL. Chemical and morphological analyses of the Lecidea lithophila-plana group (Lecideaceae), 189
CULBERSON, C. F., T. H. NASH III & A. JOHNSON. 3-α-Hydroxybarbatic acid, a new depside in chemosyndromes of some Xanthoparmeliae with β-orcinol depsides, 154
Cyathodium cavernarum, 626; indicum, 628, 637; tuberosum, 144
cyclic nucleotides, antagonism of cytokinin induced callus in Pylaisiella selwynii by, 47
Cyclodictyon albicans, 238; capillatum, 238; cuspidatum, 238; humile, 238; lindigianum, 238; nivale, 238; obscurifolium, 238; riparium, 248; roridum, 238, 248; rubrisetum, 238; varians, 238
Cynodontium bogotense, 221, 248; glaucescens, 411; rufescens, 221, 248; strictum, 221, 248; strumiferum, 496; tenellum, 411, 496
Cyphella subcyanea, 92
Cyphelium trachylioides, 320
Cyrtomnium hymenophyllum, 168, 409, 411
cytokinin induced callus in Pylaisiella selwynii, 47

Dactylina artica, 398+; beringica, 111; madreporiformis, 404; ramulosa, 404
Daltonia bilimbata, 238, 248; binervis, 238, 248; brevinervis, 238; compressa, 238, 248; contorta, 80; cucullata, 238; gracilis, 238, 248; irrorata, 238, 248; lindigiana, 238, 248; longifolia, 238; ocanniana, 238, 248; ovalis, 238; pulvinata, 238; splachnoides, 517; tenuifolia, 238; wallisii, 238, 248
daylength, effect on bryophytes, 29+
DELGADILLO M., C. Mosses and phytogeography of the Liquidambar forest of Mexico, 432
DELGADILLO M., C. Notes on alpine mosses of Mexico, 629

4-O-demethylplanaic acid, 189+
Dendriscocaulon umhausense, 404
Dendroalsia abietina, 262
Dendroceros crispus, 626
Dendroligotrichum dendroides, 519
Dermatocarpon acarosporoides, 23; compactum, 23; lachneum, 23; miniatum, 23, 25, 27; moulinsii, 23; plumbeum, 23; reticulatum, 23; tuckermanii, 23; vellerum, 23
desert mosses, desiccation of, 65+
desiccation of desert mosses, 65+
Desmatodon cernuus, 542, 557; convolutus, 80, 262, 542; guepinii, 262; latifolius, 262, 542, 519, 550; laueri, 542; leucostomus, 543; obtusifolius, 275, 485, 543, 550, 555, 609, 611; plinthobius, 262, 541, 611; porteri, 289, 543, 550; randii, 557; systylius, 543; ucrainicus, 557
development of the archegonium and embryo in Lophocolea heterophylla, 576
Dialytrichia, 546, 549
Diaphanodon procumbens, 147
Diaphodon friisii, 494
diazotrophic organisms, 55+
dichlorodepsidone, 292+
Dichodontium olympicum, 587; pellucidum, 262, 587
Dicranella angustifolia, 223; bogotensis, 223; callosa, 223; consimilis, 223; crassinervis, 248; curviseta, 248; debilis, 74; densa, 248; ditissima, 223; grevilleana, 516; heteromalla, 223, 246, 248, 275, 485, 611; hilariana, 74, 223; lindigiana, 248; macrocarpa, 224, 248; mayorii, 224; muralis, 248; palustris, 411; perrottetii, 224, 248; pusilla, 248; schreberiana, 411; strumulosa, 224, var. minor, 224; subinclinata, 224; vaginata, 248; varia, 74, 262, 275, 611
Dicranodontium denudatum, 222, 224; falcatum, 80; fleischerianum var. longisetum, 498; longisetum, 248; pusillum, 221, 248; setosum, 224
Dicranoweisia brevifolia, 587, 588; cirrata, 262, 587, 588; crispula, 516, 588
Dicranum altissimum, 222, 248; argyrocaulon, 221, 248; bolivianum, 224, 248; chionophilum, 248; chrysodictyon, 222, 248; concolor, 248; condensatum, 73, 75, 485, 492; deflexicaulon, 224; densum, 248; elongatum, 366; filicaule, 222, 248; flaccidissimum, 224, 248; flagellare, 485; frigidum, 224; fulvum, 485; fuscescens, 71; goudotii, 224; jamesonii, 248; lamprodictyon, 222, 248; longisetum, 248; macrodon, 248; montanum, 275, 609, 611; paramicola, 224; pauperum, 222, 248, var. minus, 223; polysetum, 71; porphyreodictyon, 222, 248; propinquum, 222, 248; reflexisetum, 223, 248; rosulatum, 223, 248, var. iteratum, 223; sclerocarpum, 224, 248; scoparium, 142, 485, 516, 611; speciosum, 223, 248; speirophyllum, 476; spiripes, 221, 248; spurium, 485; strictiusculum, 224; strictulum, 223, 248; subconcolor, 223, 248; vesiculare, 224; viride, 275; wallisii, 223, 248
Dictyonema interruptum, 109, 501; irpicinum forma laudatatea, 109; ligulatum, 109; pavonia, 109; sericeum, 109
Didymodon acutus, 544, 551, 554, var. ditrichoides, 499, var. icmadophila, 499; asperifolius, 413; australasii, 499; brachyphyllus, 499; constrictus, 544; fallax, 499, 544, 551, 554; giganteus, 487, 489; gracile, 224, 248; icmadophilus, 544, 551; killipii, 226; laevigatus, 499; lamyanus, 544, 551; luridus, 544, 551; mexicanus, 263; nigrescens, 487, 544, 551; occidentalis, 499; rigidicaulis, 544, 551; rigidulus, 263, 519, 539, 544, 551, 554; spadiceus, 544, 548, 551; subandreaeoides, 499; tophaceus, 263, 544, 551, 609, 611; umbrosus, 499; vinealis, 499, 544, 551, var. flaccidus, 499, 544, 551
DIETERT, M. F. Studies on the gametophyte nutrition of the cosmopolitan species Funaria hygrometrica and Weissia controversa, 417
diffractaic acid, 155+
Dimaelaena oreina, 23, 404
Dimerella bacillifera, 473; lutea, 133
Diphyscium foliosum, 29, 30, 32, 33, 275, 438, 485
Diplasiolejeunea auriculata, 637; comorensis, 637; ensifera, 637; hamata, 637; integerrima, 637; onraedtii, 310; plicatiloba, 634; ramicola, 637; tridentata, 637
Diplochistes actinostomus, 23, 125, 133; scruposus, 23
Diplophyllum albicans, 142, 145; apiculatum, 483, 609, 610; taxifolium, 167
Dissodon scabrisetus, 248
Distichium capillaceum, 220, 375, 490, 516, 630
Distichophyllum elongatum, 238; flavescens, 517; pusillum, 239, 248
Ditrichum ambiguum, 501; bogotense, 221; crinale, 221; currituckí, 592; difficile, 592; gracile, 221; henryi, 592; heteromallum, 221, 590, 591, var. emodii, 590, 591; homomallum, 516; lineare, 592; montanum, 592; pallidum, 485, 592, 611; pusillum, 476, 591, 611; rufescens, 221, 248, 435; schimperi, 592; submersum, 221; subulatum, 592; tortipes, 592
DIXON, H. N. (portrait), 124*
Drepanocladus aduncus, 263, 514; exannulatus, 240; fluitans, 240; revolvens, 73, 240; uncinatus, 364; vernicosus, 82, 83
Drepanophyllum falcifolium, 248
Drummondia prorepens, 485, var. latifolia, 496
Dumortiera hirsuta, 142, 144, 150, 152, 483

Echinoplaca intercedens, 88+, 89*, 503; pellicula, 125, 133
Ectropothecium amabile, 244, 248; amphibolum, 244, 248; apiculatum, 244, 248; campanulatum, 243; globitheca, 244, 248, 250; laevifolium, 242, 248; sandwichense, 80
effect of nutritional conditions on sexual reproduction in Riccia, 37
EGAN, R. S. Recent literature on lichens—101, 105; —102, 314; —103, 500
EGUNYOMI, A. & S. O. OLARIMOYE. Mosses from Nigeria II. Collections from the Obudu Cattle Ranch, 76
embryo in Lophocolea heterophylla, development of, 576
emodin, 302+
Encalypta affinis, 454; alpina, 455; brevicolla, 169, 450+; brevipes, 380, 414, 415, 450, 456+; ciliata, 263, 289, 450+; commutata, 455; crispata, 454; daviesii, 454; fimbriata, 454; **flowersiana**, 368+*, 415; lanceolata, 454; longicolla, 162, 169, 374, 414, 450+; mutica. 374. 377. 380. 414. 415. 450. 458+; rhaptocarpa, 368+, 411, 414, 452+, 454; spathulata, 377+, 414, 450+; squarrosa, 454; streptocarpa, 368, 452+; **vittiana**, 368+, 411, 412, 415; vulgaris, 73, 263, 368+, 414, 453+
Encalyptaceae americanae exsiccatae: An annotated guide to Fasciculus I (Numbers 1–10), 150
Endocarpum pusillum, 23, 125, 133
Enterographa elaborata, 501; sorediata, 501
Entodon aurescens, 241; beyrichii, 241; brevisetus, 263; cladorrhizans, 611; complicatus, 241; compressus, 611; consanguineus, 248; cupressiformis, 241; gracilisetus, 242; jamesonii, 242, 248; longisetus, 248; lutescens, 242, 248; macropodus, 242, 485, 492; myurus, 49, 498; onustus, 248; polycarpus, 242, 248; propinquus, 242; ramosissimus, 242; schleicheri, 494; sedutrix, 275, 485, 492, 611; squarrosus, 248; subcuspidatus, 80
Entosthodon attenuatus, 263; bolanderi, 263; californicus, 263; jamesonii, 248; lindigii, 249; lutescens, 249; pseudomarginatus, 228, 249; undulatus, 249
Eoleucodon pacificus, 496
Ephemerum serratum, 289
Epilichen glaucinigennus, 502; scabrosus, 502
Epipterygium lepidopiloides, 434
Eremonotus myriocarpus, 119, 162+
Eriodon conostomus, 515
Eriopus cristata, 517; deflexus, 238; nutans, 238
Erpodium coronatum, 232; hanningtonii, 598
Erythrodontium consanguineum, 242; longisetum, 242, 251, 252, 254; squarrosum, 242, 434, 444
erythroglaucin, 302+
Erythrophyllopsis andina, 226, 259
Erythrophyllum campylocarpum, 249
Eucatogonium politum, 249
Eucladium verticillatum, 263, 285, 476, 477, 478, 544, var. recurvatum, 477
Eurhynchiella semitorta, 241
Eurhynchium hians, 73, 275, 611; oedogonium, 241; pseudopiliferum, 241; pulchellum, 73, 241, 263, 275, 375, 611; riparioides, 263; semiscabrum, 241
Evernia perfragilis, 404
Everniastrum paramense, 502
Everniopsis trulla, 110

Fabronia ciliaris, 275, 438, 442, 611; lindigiana, 240; polycarpa, 240; pusilla, 263, 517; wrightii, 434, 441
fallacinal, 302+
fallacinol, 302+
Fegatella conica, 98, 141
Felipponea montevidensis, 518
FIFE, A. J. Taxonomic observations on three species of North American Funaria, 204
Fissidens **abruptus**, 58+, 59*; angustifolius, 58, 60, 61, 267+, 479; asplenioides, 220, 268, 438; brittonii, 58, 61, var. percurrens, 61; borealis, 496; bryoides, 63, 74, 266+, 275, 438, 610, 611, 630; bushii, 289+, 611; camptodontius, 220; cristatus, 275, 426, 435, 438, 443, 611; densiretis, 479; diplodus, 435; dissitifolius, 267, 436, 618; divisus, 220; donnellii, 436, 441; elegans, 220, 268+, 436, 444; ensifolius, 58, 60; excurrentinervis, 268+; exiguus, 610; exilis, 517; flavinervis, 220; flexinervis, 249; fontanus, 275, 438, 485; garberi, 268; grandifrons, 263, 485; hancockianus, 220; hyalinus, 267+; intermedius, 80; intromarginatus, 220; kegelianus, 61, 63, 64, 268; limbatus, 263; lindigii, 220, 249; leptopodus, 268; longidecurrens, 268; macroblastus, 58, 61, 64; milobakeri, 58, 61, 64, 266; minutulus, 485; mollis, 266+, 479; Nomenclatural and distributional notes on some American species, 58; obscurocostatus, 60; obtusifolius, 273+, 485, 611; osmundoides, 290, 485; papulans, 58, 60; polypodioides, 220, 437; prionodes, 220, 249; pusillissimus, 62; radicans, 62, 479; rambii, 144; repandus, 220, 249, 436, 441; rigidulus, 58, 62, 220, 269; steerei, 58, 62, 479, 618; subbasilaris, 60, 437, 444, 485; svihlae, 58, 60; taxifolius, 275, 438, 485, 611; tequendamensis, 220, 249; **termitarum**, 58+, 59*; tortilis, 249; vardei, 436; viridulus, 275, 610; wallisii, 220
Florschuetziella steerei, 1+, 13*, 15*, 17*
FLORSCHÜTZ-DE WAARD, J. & P. A. FLORSCHÜTZ. Estudios sobre criptógamas colombianas III. Lista comentada de los musgos de Colombia, 215

Foliicolous lichens from southeastern United States, 88
Folioceros buitenzorgius, 632; dixitianus, 632; fuciformis, 632; fuscus, 632; glandulosus, 632; incurvus, 632; indicus, 632; mamillisporus, 632; mangaloreus, 632; miyabenus, 632; physocladus, 632; pinnilobus, 632; spinisporus, 632; vesiculosus, 632; weistii, 632
Fontinalis antipyretica, 263, 382+, 594+; bogotensis, 233; duriaei, 382+; hypnoides var. plicatus, 496; novae-angliae, 383, 391, 492; squamosa, 389, 594+
Forssellia umbilicatá, 506
Forsstroemia trichomitria, 485, 492
Fossombronia angulosa, 112, 634; brasiliensis, 299, 483; foveolata, 299
fragilin, 302+
FRAHM, J.-P. New Campylopus records from Mexico, including C. sharpii sp. nov., 623
FRECKMAN, S. *See* M. COLE, F. D. BOWERS & S. FRECKMAN, 273
FREEMAN, G. W. *See* L. M. O'FLAHERTY & G. W. FREEMAN, 609
Frullania apiculata var. obtusata, 115; brittoniae, 310; curvistipula var. latistipula, 311; dilatata, 114, 116; eboracensis, 274, 299, 483, 610; evelynae, 311; eymae var. dentistipula, 311; fragilifolia, 114; gemmipara, 311; gracilis subsp. zennoskei, 311; honostipula, 634; inflata, 299, var. dioica, 311; koponenii, 116; neosheana, 634; nobilis, 311; pachyderma, 311; pariharii, 115, 311; retusa var. gymantha, 115, var. hirsuta, 311; riparia, 299, 483; sheana, 634; spinigastria, 634; squarrosa, 299, 483; subdentata, 311; subvalida, 115; tamarisci subsp. asagrayana, 274, 483, subsp. obscura, 311; tuyamae, 115
Fulgensia bracteata, 302+; fulgens, 302+, 504; fulgida, 303; subbracteata, 303, 504
fulgidin, 302+
fulgoicin, 302+
Funaria arctica, 204+, 208*; bonplandii, 228, 436; calvescens, 206, 228, 249; flavicans, 73, 74, 205+, 421, 485, 611; **groutiana**, 204+, 208*; hygrometrica, 47, 50, 52, 96, 144, 205+, 228, 263, 275, 417+, 485, 517, 540, 611, var. arctica, 207, var. calvescens, 228, 249, 438, var. minor, 213; jamesonii, 228; lindigii, 228; lutescens, 228; microstoma, 74, 204+, 208*, var. obtusifolia, 204+, forma canadense, 205+; muhlenbergii, 263; obtusifolia, 207+; submicrostoma, 205; trumffii, 228, 246, 248; undulata, 228
Fuscidea cyathoides var. sorediata, 318; intercincta, 318; lightfootii, 315

gametophyte nutrition in Funaria hygrometrica and Weissia controversa, 417
gangaleoidin, 292+
Garckea phascoides, 588, 593
Geocalyx graveolens, 114
geographic disjunction and heterophylly in Tortella fragilis var. tortelloides (=Sarconeurum tortelloides), 84
Geothallus tuberosus, 98
Gigaspermum repens, 598, 599
GILBERT, O. L. *See* M. M. MAHANDRU & O. L. GILBERT, 292, 302
GLIME, J. M. & D. W. ACTON. Temperature effects on assimilation and respiration in the Fontinalis duriaei-periphyton association, 382
Globulinella peruviana, 630
Glossadelphus chrysobasilaris, 80; laevifolius, 242; truncatulus, 242
Glyphis cicatricosa forma confluens, 473
Glyphocarpa laevisphaera, 249; lindigii, 231, 249; strumosa, 249; taylorii, 230, 249
Glypholecia scabra, 23
Glyphomitrium daviesii, 454; lindigii, 249; lobuliferum, 232, 249
GODFREY, J. D. & W. B. SCHOFIELD. New and interesting hepatics from British Columbia, Canada, and northern Washington state, U.S.A., II., 162
Gomphillus americanus, 315
Goniobryum subbasilare, 520
Gram-negative bacteria, 141
Gram-positive bacteria, 141
Graphina carneoviridis, 320; columbiana, 320; illinata, 320; nylanderiana, 473; plurispora, 320; scolecitis, 125, 133; suberythrella, 320; triphoroides, 320
Graphis afzelii, 133; caesiella, 134, 473; imshaugii, 320; isidiifera, 320; scripta, 127, 473
GRIFFIN, D. III. New records for the moss flora of Venezuela, 618
Grimmia affinis, 227; agassizii, 279; alpicola, 485, var. dupretii, 263, var. rivularis, 279; anodon, 67+, 263, 609, 610, 611; apocarpa, 263, 611, 622, var. conferta, 263, var. pulvinata, 80, var. stricta, 79, 609, 610; bogotensis, 227; calyptrata, 263; cinclidodontea, 263, 279+; contermina, 227, 249; cribrosa, 411, 412, 415; crispipila, 227, 249; cucullatifolia, 249; doniana, 596, 597; hartmanii, 263; kiriensis, 496; laevigata, 73, 80, 263, 485, 492; lanuginosa, 249; maritima, 279; montana, 263; moxleyi, 263; occidentalis, 279+; orbicularis, 67+; ovalis, 227, 249, 263; ovata, 227, 249; **pacifica**, 276+, 278*; pilifera, 485; plagiopoda, 73, 517; pulvinata, 67+, 263; tenerrima, 263; trichophylla, 263; unicolor, 74
Groutiella apiculata, 1+, 10*, 249, 256, 436; chimborazense, 436, subsp. chimborazense, 3*+, subsp. **reesei**, 1+, 3*, 17*; fragilis, 232, 249; mucronifolia, 1+, 232, 249, 251,

256, 480; tomentosa, 232, 249, 259, 440, 443; **tumidula**, 1+, 10*; undosa, 1+; wagneriana, 1+
growth, reproduction and population ecology in relation to microclimate in the bipolar moss Polytrichum alpestre, 325
Guembelia bogotensis, 227, 249
Gyalecta foveolaris, 404
Gyalectidium filicinum, 88, 91; rotuliforme, 88, 91, 110
Gyalidia diaphana, 506
Gymnomitrion apiculatum, 162+; concinnum, 164*; corallioides, 168
Gymnostomiella orcuttii, 619
Gymnostomum aeruginosum, 85, 274+, 485, 544, 551, 554, 609, 611; bonplandii, 249; calcareum, 263, 519, 544, 551, 552, 622; recurvirostrum, 264, 275, 485, 611, var. latifolium, 485
Gyroweisia lindigii, 226; nigeriana, 495; reflexa, 544; tenuis, 544, 554, var. badia, 544

habitat ecology of Tasmanian lichens, 171+
Haematomma caesium, 315; lapponicum, 404; ochroleucum, 317, var. porphyrium, 317
Haplocladium macropilum, 518; microphyllum, 264, 438, 485, 611; virginianum, 73, 275, 485, 609, 611
Haplodontium jamesonii, 229, 247, 249; megalocarpum, 229, 249
Haplohymenium triste, 80, 275, 439, 444, 485
Haplolejeunea cucullata, 634
Haplomitrium gibbsiae, 579; grollei, 634; hookeri, 115; mnioides, 115
HARRIS, C. W. (portrait), 324*
HARRIS, R. C. Four species of Thelopsis Nyl. (lichenized Ascomycetes) new to North America, 77
Harrisonia humboldtii, 249; rhabdocarpa, 234, 249
HASSE, H. E. (portrait), 324*
HARTHILL, M. P., D. M. LONG & B. D. MISHLER. Preliminary list of southern California mosses, 260
Hawaiian mosses, new and noteworthy records for, 79
Hedwigia ciliata, 233, 249, 264, 485, 497, 611; humboldtii, 234, 249; imberbis, 234, 249; subrevoluta, 249, var. incana, 233
Hedwigidium integrifolium, 234, 249
HEGEWALD, E. *See* R. A. PURSELL & E. HEGEWALD, 267
Helicoblepharum daltoniaceum, 238, 248
Helicodontium capillare, 436
Helicophyllum torquatum, 233, 436, 619
Helodium blandowii, 82, 83
Hemiragis aurea, 238, 619
Heppia lutosa, 23
Herbertus aduncus, 166; sakurai subsp. arcticus, 312; stramineus, 637
Herpetineuron toccoae, 439, 442
Herpothallon sanguineum, 139
HERRE, A. W. C. T. (portrait), 324*
HERTEL, H. *See* C. F. CULBERSON & H. HERTEL, 189
Herzogiella seligeri, 414; turfacea, 414
Heterocladium dimorphum, 414
Heterophyllium affine, 249, 439; nemorosum, 242, 249, 250, 255
Heterophylly in Tortella fragilis var. tortelloides (=Sarconeurum tortelloides), 84
Heterothecium augustini, 92; puiggarii, 92
HOE, W. J. Additional new and noteworthy records for Hawaiian mosses—5, 79
HOE, W. J. & R. H. ZANDER. Geographic disjunction and heterophylly in Tortella fragilis var. tortelloides (Sarconeurum tortelloides), 84
HOLCOMBE, J. W. Sphagnum angermanicum in New Hampshire, 616
Holomitrium arboreum, 436; crispulum, 224; flexuosum, 434; lehmannii, 224; longifolium, 224; seticalycinum, 80; urvilleanum, 497
HOLZAPFEL, W. H. *See* D. C. J. WESSELS, L. A. WESSELS & W. H. HOLZAPFEL, 270
HOLZINGER, J. M. (portrait), 324*
Homalia glabella, 436, 480; trichomanoides, 485
Homaliodendron decompositum, 236, 436
Homalotheciella subcapillata, 485
Homalothecium aeneum, 264; arenarium, 264; leskeoides, 241, 250, 251, 255, 439; nevadense, 264; nuttallii, 264; pinnatifidum, 264
Homomallium adnatum, 73, 275, 485, 611
HONG, W. C. The genus Scapania in western North America I. Historical background, 181
Hookeria albicans, 249; bogotensis, 237, 249; capillata, 238, 249; crispa, 238, 249; crugeriana, 249; cuspidata, 249; diversifolia, 249; falcata, 238, 249; gemmacea, 237, 249; glandulifera, 238, 249; hispidula, 237, 249; humilis, 238, 249; hypnacea, 249; leiophylla, 517; lindigiana, 238, 249; nivalis, 249; ovata, 237, 249; pallida, 249; pilifera, 249; quadrifaria, 517; rorida, 238, 249; rubriseta, 238, 249; subfalcata, 239, 249; taylorii, 249; variabilis, 239, 249; velutina, 239, 249; viridissima, 239, 249
Hookeriopsis acuminata, 238; crispa, 238; cruegeriana, 238, 436; diffusa, 238; falcata, 238; glandulifera, 238; gracilis, 238; heteroica, 433+; hypnacea, 238; incurva, 480; killipii, 239, 249; negrensis, 239; pendula, 249; subfalcata, 239; taylorii, 239, undata, 239; variabilis, 239; velutina, 239; viridissima, 239
HOOPER, G. R. *See* K. K. BAKER, J. A. MALACHOWSKI & G. R. HOOPER, 533

HORTON, D. G. Encalypta vittiana sp. nov. and E. flowersiana sp. nov. from North America, 368
HORTON, D. G. *See* D. H. VITT & D. G. HORTON, 409
HORTON, D. G. Encalyptaceae americanae exsiccatae: An annotated guide to Fasciculus I (Numbers 1–10), 450
Huilia crustulata, 396+; flavocaerulescens, 404; glaucophaea, 404; macrocarpa, 404; musiva, 506, var. subsquamescens, 506; panaeola, 404
Husnotiella revoluta, 264
Hydrogonium arcuatum, 144, 544; javanicum, 544; subcomosum, 544
3-α-Hydroxybarbatic acid, a new depside in chemosyndromes of some Xanthoparmeliae with β-orcinol depsides, 154
Hygroamblystegium fluviatile, 275, 611; irriguum, 514; tenax, 264, 611
Hygrobiella laxifolia, 163
Hygrohypnum closteri, 290+; dilatatum, 290; molle, 264, 289+; ochraceum, 290; pelichucense, 240, 244; steerei, 498; tequendamense, 240, var. subfalcatum, 240
Hylocomium pyrenaicum, 410; splendens, 328, 361, 366, 367, 431
Hymenostomum microstomum, 520
Hymenostylium recurvirostrum, 545, 554, var. aurantiacum, 545
Hypnella pilifera, 239
Hyophila acutifolia, 544; involuta, 147, 150, 273+, 476, 479, 485, 544, 560, 609, 610; lindigii, 226, 249; propagulifera, 544; tortula, 226, 437
Hypnum acrophyllum, 239, 250; affine, 242, 243, 250; agnatum, 243, 250; amabile, 244; andinum, 242, 250; anomalum, 235, 250; apiculatum, 244, 250; aquaticum, 241, 250; arzobispoae, 240, 250; asperulum, 240, 250; aureo-sulfureum, 243, 250; bonplandii, 250; caespitosum, 243, 250; carantae, 240, 250; chrysocladon, 240, 250; chrysophylloides, 241, 250; chrysostegum, 250; clinocarpum, 241, 250; conchophyllum, 250; crassiusculum, 250; crithmifolium, 243, 250; cucullatifolium, 243, 250; cupressiforme, 74, 244, 497, 517, 539, 557; curvifolium, 485; delicatulum, 240, 250; elatulum, 236, 250; exasperatum, 241, 250; fasciculatum, 250; flagelliferum, 236, 250; fragillirostrum, 243, 250; frontinoae, 240, 250; globitheca, 250; gratum, 250; hamatum, 244; hians, 498; hispidulum, 250; hochstetteri, 496; hospitans, 242, 250; imponens, 485; intermedium, 250; kegelianum, 250; laevifolium, 241, 250; lancifrons, 236, 250; langsdorffii, 244, 250; lindbergii, 74, 275, 485, 611; lindigii, 243, 250; loriforme, 244, 250; loxense, 243, 250; muricatulum, 250; neckeraeforme, 236, 250; nigricans, 235, 250; novo-granatense, 242, 250; occidentale, 241, 250; pallescens, 275; patens, 235, 250; pauperum, 240, 250; perspicuum, 244, 250, var. molle, 244, var. teres, 244; phyllogonium, 250; plumaeforme, 147; plumosum, 250; poeppigianum, 250; praelongum, 241, 250, 498; prasophyllum, 241, 250; pseudo-delicatulum, 240, 250; pseudo-lutescens, 241, 250; pseudo-piliferum, 241, 250; pseudo-protensum, 250; pseudo-rutabulum, 241, 250; pygmaeum, 250; rugisetum, 241, 250; scariosum, 251; schlimii, 251; schreberi, 251; semitortum, 241, 251; solutum, 251; sparsifolium, 236, 251; sparsum, 236, 251; spiripes, 244; sticticola, 243, 251; subbrachycarpum, 243, 251; subimponens, 244, 251; subnitidum, 243, 251; subobscurum, 244, 251; subrotundum, 241, 251; subscabrum, 251; tamariscinum, 251; tequendamense, 251; thelistegum, 244, 251; tocaremae, 241, 251; toxarion, 251; triquetrum, 36; ulicon, 251; vaucheri, 74; viscidulum, 244, 251; volvatum, 244
Hypogymnia austerodes, 404; enteromorpha, 280+, 564−; lugubris, 174; oroarctica, 404; physodes, 180, 315, 316, 404, 571; subobscura, 404
Hypopterygium rigidulum, 239; scutellatum, 239, 251; tamariscinum, 239, 251, 439, 480
Hypotrachyna cendensis, 502; meridensis, 502; neoflavida, 502; primitiva, 502

Icmadophila ericetorum, 404
Illecebraria julacea, 251
Illinois, eleven mosses new to, 609; phenology of selected bryophytes in southern, 29
impact of emissions from coal-fired electricity generating facilities on N_2-fixing lichens, 54
India, Notothylas dissecta, a hornwort new to, 625
induced callus in Pylaisiella selwynii, 47
Ionaspis epulotica, 404; melanocarpa, 404; ochracella, 404
IRELAND, R. R., JR. *See* R. H. ZANDER & R. R. IRELAND, JR., 474
irradiance values, 30+
Isodrepanium lentulum, 239, 436, 480
Isopterygiopsis muelleriana, 412, 609, 611
Isopterygium chrismarii, 242; deplanatiforme, 242; diminutivum, 242, 434; distichaceum, 275, 291; elegans, 79, 290, 411; hospitans, 242; planissimum, 517; pseudo-tenerum, 242; pulchellum, 275; tenerum, 242, 250
Isothecium cristatum, 264; spiculiferum, 264
Iwatsukiella leucotricha, 494

Jaegerina scariosa, 560
JACOBSON, D. E. & P. V. PRIOR. Additions to the moss flora of Nebraska, 622

JOHNSON, A. *See* C. F. CULBERSON, T. H. NASH III & A. JOHNSON, 154
JOURNET, A. R. P. (review), J. R. Etherington, Plant physiological ecology, 508
Jubula hattori, 637; pennsylvanica, 483
Jungermannia acuta, 581; atrovirens, 118; hyalina, 274; infusca, 635; nemorea, 185; nemorosa, 185; rosulans, 635; sphaerocarpa, 117; thermarum, 635, torticalyx, 117, 635; tristis, 118, 635

KANDA, I. *See* N. KITAGAWA & I. KANDA, 96
Kingiobryum paramicola, 224
KITAGAWA, N. & I. KANDA. Linear spore tetrads in Conocephalum supradecompositum, 96
KUMAR, N. R. *See* K. R. RAO, N. R. KUMAR & A. N. REDDY, 286
KUMAR, S. S. & S. K. VERMA. Cytological investigations on some West Himalayan mosses, 583
KUWAHARA, Y. Notes on the genus Steereella, 613
Kurzia touwii, 116

LANGE, O. L. *See* P. W. RUNDEL, G. C. BRATT & O. L. LANGE, 171
Lasallia papulosa, 23; pensylvanica, 404; pustulata, 314
LAWTON, E. Grimmia pacifica, a new species from western North America, 276
Lecanora alphoplaca, 23, 25, 27; atra, 23; badia, 404; carpinea, 23; castanea, 404; cenisia, 23; chalcophila, 506; charona, 23; chlarotera, 23, 292+; chrysoleuca, 24, 564; coilocarpa, 404; conizaeoides, 25, 318; crenulata, 23; dispersa, 23, 404; elevata, 396+; epibryon, 404; frustulosa, 23; gangaleoides, 292+, 502; garovaglii, 23; hageni, 23, 396+; margacea, 318; melanophthalma, 21, 27; muralis, 23, 25; mutabilis, 23; nordenskioeldii, 404; novomexicana, 23, 27; piniperda, 23; polytropa, 404; rugosella, 396+; rupicola, 23, 404; saligna, 23; sulphurea, 292; vegae, 396+
Lecidea aglaea, 396+; antoniensis, 191+; armenica, 404; assimilata, 404; atroalba var. enteromorpha, 196; atrobrunnea, 23, 404; auriculata, 404; berengeriana, 404; cinerorufa, 404; confluens, 404; decipiens, 23; elegantior, 405; farinosa, 196; fusca, 405; fuscoatra, 197; globifera, 23; granulosa, 23; hassei, 191+; heteromorpha, 196; hypocrita, 405; lactea, 405; lapicida, 405; leptoboloides, 189+; lichenicola, 502; lithophila, 189+; lithophiloides, 192+; lithophilopsis, 189+; lucida, 405; macrocarpa, 197; plana, 189+; rubiformis, 405; russellii, 23; symmicta, 405; tessellata, 23, 405; tornoensis, 405; turgidula, 23; vernalis, 405; vorticosa, 405
Lecidella carpathica, 23; glomerulosa, 23, 405; stigmatea, 23; viridans, 23
Leersia fimbriata, 453; lanceolata, 453; vulgaris, 453
Leiocolea alpestris, 115; collaris, 115, 118; gillmannii, 118; kateninir, 118
Leiomela aristata, 231, 251; aristifolia, 231, 251; bartramioides, 231, 246, 248, 439, 476; lopezii, 231, 257; lutescens, 231
Leiomitrium scaberrimum, 1, 16; standleyi, 1, 16
Lejeunea alaskana, 312; magohukui, 115; mizutanii, 634
Lepidolejeunea radicosa, 634
Lepidopilum angustifrons, 239, 251; apollinairei, 239; aubertii, 239; biductulosum, 239, 251; brevipes, 239; carneum, 239; caudatum, 239; daltoniaceum, 238, 251; erubescens, 239; excelsum, 239; flexifolium, 239, 251; frondosum, 239; hyalinum, 251, 349; longifolium, 239; maculatum, 239; mniaceum, 239; muelleri, 239; nutans, 239, 251; pergracile, 239; perlaxum, 239; permarginatum, 239; polytrichoides, 239, 480; pringlei, 436; purpurascens, 619; radicale, 239, 436; robustum, 239; scabrisetum, 239, 251; sprucei, 251; subenerve, 251; tortifolium, 239; wallisii, 239
Lepidozia parvula, 116; reptans, 274; vitrea, 117
Lepraria incana, 23
Leptobarbula verica, 544, var. winteri, 544
Leptobryum pyriforme, 229, 264, 274, 611
Leptochlaena graciliseta, 251
Leptodictyum brevipes, 609, 610; riparium, 264, 275, 480, 525, 540, 611; trichopodium, 264, 275, 611
Leptodon smithii, 518
Leptodontium acutifolium, 226, 251; brachyphyllum, 227; capituligerum, 226, 251; erythroneuron, 226; filescens, 226, 251; filicola, 226; flexifolium, 226, 251, 255, 545, 554; fuhrmannii, 226; gracile, 226, 251; gracilescens, 226, 251; longicaule, 226, 251; luteum, 226; orthotrichoides, 226; planifolium, 226; proliferum, 226; pungens, 226, 251; spinosum, 233, 251; squarrosum, 520; stoloniferum, 226; subgracile, 251; subgrimmioides, 226, 251; sulphureum, 227, 251; syntrichioides, 227; ulocalyx, 227, 251; vaginatum, 227, 251; viticulosioides, 227, 454, var. exasperatum, 227, var. panamense, 227, 251, 252, 256, 437, 443, var. viticulosoides, 439, 444; wallisii, 227, 251
Leptogium austroamericanum, 473; cyanescens, 473; lichenoides, 405; millegranum, 124, 134; phyllocarpum, 473; saturninum, 405
Leptohymenium consanguineum, 242, 251; cylindricaule, 242, 251, var. rupestre, 242;

longisetum, 242, 251; squarrosum, 242, 251; tenue, 434
Leptostomum inclinans, 518
Leptotheca costaricensis, 230
Leptogium africanum, 108; chiriquiense, 108; glaucescens, 134; hypoprotocetraricum, 108; lepadodes, 125, 134; mammiculum, 108; panamense, 108; polycarpum, 125, 134; santense, 134; stellatum, 108; wightii, 134
Leptotrichum rufescens, 251
Lepyrodon suborthostichus, 234, 251; tomentosus, 234, 251, 436, 443, 444
Leskea angustata, 240, 251; bonplandii, 241, 251; gracilescens, 275, 611; gracillima, 240, 251; involvens, 251; obscura, 275, 611; polycarpa, 275, 518, 611
Leskeodon andicola, 239; pusillus, 239; wallisii, 239
Letharia vulpina, 111
Leucobryum albidum, 439, 485; antillarum, 225, 436, 441; crispum, 216; giganteum, 216, 225; glaucum, 225, 251, 275, 485, 609, 611; martianum, 225, 436; obtusifolium, 225, 251; polakowskyi, 225, 436; vulgare, 251
Leucodontopsis geniculata, 234, 560
Leucoloma cruegerianum, 436; macrodon, 224; procumbens, 224; scaberulum, 81; serrulatum, 224, 436; subimmarginatum, 436, 444
Leucodon julaceus, 485; macrosporus, 434, 441
Leucolejeunea clypeata, 423, 431, 484
Leocomium acrophyllum, 239; compressum, 239; lignicola, 239
lichen-feeding Coleoptera, 270
lichen-invertebrate associations, 270
lichenized Ascomycetes, 77
Lichenodiplis lichenicola, 502
lichens from Louisiana, new or noteworthy records of, 125
lichens in southern Ohio, cold injury to, 620
lichens, recent literature on, 105, 314, 500
Lindbergia brachyptera, 73, 275, 290, 485, 611
Lindigia aciculata, 235, 250, 251, 436; curtipes, 251; debilis, 235; hypnoides, 235, 251; imponderosa, 235, 251; pilotrichelloides, 235, 251
linear spore tetrads in Conocephalum supradecompositum, 96
lipids of the sperm masses of three mosses, 93
Lobaria gyrophorica, 317; linita, 405; orientalis, 317; pulmonaria, 319; scrobiculata, 396*; yoshimurae, 317
LONG, D. M. *See* M. P. HARTHILL, D. M. LONG & B. D. MISHLER, 260
LONGTON, R. E. Studies on growth, reproduction and population ecology in relation to microclimate in the bipolar moss Polytrichum alpestre, 325
Lopadium augustini, 92; flammeum, 93; fuscum, 88, 92, 125, 134, 505; pezizoideum, 405; puiggarii, 88+, 89*, 125, 134, 505; subcoerulescens, 92
Lophocolea alata, 118; bidentata, 118, 119, 484, 581, var. latifolia, 119; cuspidata, 118, 484, 610; heterophylla, 29, 30, 32, 86, 87, 484, 576+, 610; minor, 610
Lophozia collaris, 581; ehrhartiana, 312; gillmanii, 163; holmenianum, 312; lycopodioides, 167; obtusa, 118; quadriloba, 166+; schusterana, 118; sudetica var. anomala, 118
Lopidium concinnum, 518
Louisiana, new or noteworthy records of lichens from, 125
Lunularia cruciata, 116, 148, 633
Lyellia crispa, 143, 144

MACOUN, J. (portrait), 511*
Macrocoma filiforme, 519; frigida, 1, 14, 15, 232; intricata, 80; orthotrichoides, 435; **scaberrima**, 1, 16; sullivantii, 437
Macrodiplophyllum microdontum, 412
Macromitrium aciculare, 519; altituberculosum, 12, 435; apiculatum, 232, 251; attenuatum, 232; aureum, 232; brachyrhynchum, 1, 6, 7; brevisetum, 80; chimborazense, 8; cirrosum, 232, 436, 480; contextum, 1, 7, 232, 251; crenulatum, 232; didymodon, 251; fragilicuspis, 435; frigidum, 14, 232, 252; guatemalense, 232, 252, 435; hirtellum, 6; incurvum, 497; leprieurii, 6; liberum, 232, 252; longifolium, 232; longirostre, 232; macrostomum, 12, 14; mucronifolium, 232, 252; orthotrichoides, 14, 15; osculatianum, 216; pellucidum, 12, 216; pentagonum, 6; pentastichum, 1, 6, 7; podocarpi, 5, 6, 232; portoricense, 5, 6, 232; pringlei, 14; proliferum, 232; punctatum, 1, 6, 7, 232, 436; ramosissimum, 227, 252; reflexifolium, 7; richardii, 5, 6, 7, 232, 251, 436; schwaneckeanum, 619; **sharpii**, 1+, 3*; serrulatum, 232; stellulatum, 11; stratosum, 5, 6, 436; sullivantii, 14, 15; sumichrastii, 7; swainsonii, 233; tocaremae, 233; trachypodium, 233; trianae, 232, 252; tumidulum, 9, 11, 14; undosum, 8; undulatum, 227, 252; wagnerianum, 9
MAGILL, R. E. *See* D. M. J. MUELLER & R. E. MAGILL, 298
MAHANDRU, M. M. & O. L. GILBERT. Chemical studies in Fulgensia: structures of two new chlorodepsidones, 302
MAHANDRU, M. M. & O. L. GILBERT. Norgangaleoidin, a dichlorodepsidone from Lecanora chlarotera, 292
MAHU, M. Familias y géneros de musgos Chilenos, 513

MALACHOWSKI, J. A. *See* K. K. BAKER, J. A. MALACHOWSKI & G. R. HOOPER, 533
Malagasy Cololejeunea, some new species of, 602
Mannia californica, 484; fragrans, 112, 610; triandra, 484
MANUEL, M. G. (review), J. Maschke, Taxonomische Revision der Laubmoosgattung Myurium (Pterobryaceae), 120
Marchantia domingensis, 300; nepalensis, 148; paleacea, 114, 141+; palmata, 145, 148, 151, 286+; polymorpha, 114, 116, 117, 118, 141, 144, 210, 274, 286+, 423, 429, 431, 484, 610, 633, 635, 637; stellata, 141
Maronea constans, 125, 134
MARSH, J. E. & T. H. NASH III. Lichens in relation to the Four Corners power plant in New Mexico, 20
Marsupella condensata, 162+; revoluta, 409+, 457, 467; sprucei, 118; ustulata forma sprucei, 118
Martinellia spitsbergensis, 185
Masonhalea richardsonii, 397+
Massula elongata, 118
MATTERI, C. M. Notas briológicas (Musci) IV y V, 99
Meesia hexasticha, 414; larvardei, 230; longiseta, 230
Melanotheca cruenta, 134
Melaspilea tribuloides, 134
Menagazzia aenofusca, 174
Mesochaete crenulata, 306
Mesonodon onustus, 242
Metacalypogeia schusterana, 637
Meteoridium remotifolium, 235, 252, 253
Meteoriopsis consimilis, 235, 252; patens, 252; patula, 252, 439, 480, 560; reclinata, 144; recurvifolia, 252; remotifolia, 235, 252, 439
Meteorium appressum, 235, 252; atroluteum, 235; consimile, 235, 252; deppei, 252; excavatum, 234, 252; filiferum, 252; flexile, 252; fusco-viride, 235, 252; illecebrum, 235, 252, 436; imponderosum, 252; leucotrichum, 252; longe-barbatum, 252; nigrescens, 252; nigricans, 252; patens, 252; remotifolium, 235, 252; tenuissimum, 252; teres, 435; ternstroemii, 252; viride, 252
Metzgeria conjugata, 117, 484; cylindra, 311; decipiens, 116; foliicola, 116; fukuokana, 311; furcata, 582; japonica, 311; laciniata, 311; minor, 311
Mexico, mosses and phytogeography of the Liquidambar forest, 432; new Campylopus records from, including C. sharpii sp. nov., 623; new taxa and new combinations in the Orthotrichaceae, 1; notes on alpine mosses of, 629
Microcalicium ahlneri, 111; arenarium, 111; conversum, 111; subpedicellatum, 111
microclimate, 325+
Microdus densus, 224; lindigianus, 334; macrocarpus, 224; muralis, 224; pusillus, 224
Microlepidozia pauciflora, 118
Microthamnium andicola, 252; langsdorffii, 252; loriforme, 252; pelichucense, 244; reptans, 252; smithii, 244; substriatum, 244, 252; thelistegum, 244, 252; tuerckheimii, 244, 252; viscidulum, 252; volvatum, 252
Mielichhoferia antisanensis, 229; bogotensis, 229; brevicaulis, 229; campylocarpa, 229; clavellata, 229; dolichotheca, 229; elegans, 229; graciliseta, 229; lindigii, 229; macrocarpa, 489; megalocarpa, 229, 252; microstoma, 229, 252; mielichhoferii, 411, 412; pectinata, 229; pusilla, 229, 252; recurvifolia, 476, 477; subobliqua, 229
MILLER, N. G. (review), W. C. STEERE & H. A. CRUM, New combinations and new taxa of mosses proposed by Nils Conrad Kindberg, 507
MISHLER, B. D. *See* M. P. HARTHILL, D. M. LONG & B. D. MISHLER, 260
Mittenothamnium andicola, 244; diminutivum, 244, 250, 251, 252, 255, 439, 481, var. molle, 244, var. teres, 244; langsdorffii, 244; lehmannii, 244, 481; loriforme, 244; reptans, 244, 252, 436; squarrosulum, 434; subobscurum, 244; substriatum, 244; viscidulum, 244; volvatum, 244, var. gracilius, 144
Mniadelphus wallisii, 239, 252
Mniobryum carneum, 515; wahlenbergii, 229, 247, 619
Mniomallia viridis, 480
Mnium affine, 93+, 275; ciliare, 485, 611; cuspidatum, 142, 157, 275, 485, 611; drummondii, 275; hornum, 367; hymenophyllum, 415; lindigii, 230, 252; longirostrum, 143, 146; marginatum, 264, 275, 485, 609, 611; polycarpum, 230, 252; punctatum, 275, 486, var. elatum, 83; rostratum, 230, 252, 264, 438, var. ligulatum, 230, var. occidentale, 230; rugicum, 611; spiniforme, 252; stellare, 486; thomsonii, 275, 609, 610; undulatum, 361, 367
Moenkemeyera abrupta, 60; termitarum, 62
MOHR, C. T. (portrait), 124*
Molendoa hornschuchiana, 487+, 520, 545; sendtneriana, 85, 227, 414, 439, 488, 489, 545
Monoselenium tenerum, 116
Montana, impact of emissions from coal-fired electricity generating facilities on N_2-fixing lichens, 54+
Morinia crassicuspis, 86; ehrenbergiana, 227
MOSER, T. J., T. H. NASH III & J. W. THOMSON. Lichens of Anaktuvuk Pass, Alaska, with emphasis on the impact of caribou grazing, 393
moss flora of Nebraska, additions to, 622

mosses from Nigeria II. Collections from the Obudu cattle ranch, 76
mosses of the Great Plains III. Additions to Nebraska and the Black Hills of South Dakota and Wyoming, 72
MUELLER, D. M. J. (review), C. Suire (editor), Congrès International de Bryologie, 122
MUELLER, D. M. J. (review), R. M. SCHUSTER, Boreal Hepaticae, A manual of the liverworts of Minnesota and adjacent regions, 322
MUELLER, D. M. J. (review), M. D. GUIRY, A concensus and bibliography of Irish seaweeds, 509
MUELLER, D. M. J. & R. E. MAGILL. Notes on Texas hepatics, 298
Muelleriella crassifolia, 519
Muscoflorschuetzia pilmaiquen, 494, 515
Mycoblastus sanguinarius, 405
Mycoporellum stigmatophorum, 505
Mycoporum hypocastani, 501; quercus var. ilicis, 501
Mylia taylorii, 117; verrucosa, 117
Myurella sibirica, 275, 486, 489; tenerrima, 264
Myurium doii, 496; hebridarum, 496
Myuroclada maximowiczii, 488

NASH, T. H. III. *See* T. J. MOSER, T. H. NASH III & J. W. THOMSON, 393
NASH, T. H. III. *See* C. F. CULBERSON, T. H. NASH III & A. JOHNSON, 154
NASH, T. H. III & J. E. MARSH. Lichens in relation to the Four Corners power plant in New Mexico, 20
NASH, T. H. III & L. L. SIGAL. Gross photosynthetic response of lichens to short-term ozone fumigations, 280
Nebraska, additions to the moss flora of, 72, 622
Neckera atro-lutea, 235, 252; bogotensis, 236; chilensis, 236; consimilis, 235, 252; cylindricaulis, 242, 252; densa, 252; disticha, 252; filifera, 252; flabellata, 237; fusco-viridis, 235, 252; jamesonii, 236; lindigii, 236; longebarbata, 252; longiseta, 242, 252; luteovirens, 252; moritzii, 236, 252; nigricans, 252; novae-granadae, 236; obtusifolia, 252; pachygaster, 252; pennata, 74, 518; remotifolia, 235, 252; rufescens, 235, 253; substriata, 236, 253; ternstroemiae, 253; turgescens, 235, 253; urnigera, 435, 444
Neckeradelphus obtusifolius, 236
Neckeropsis disticha, 236, 560; undulata, 236, 436, 480, 560
Neobryum costatum, 229, 253
Neohypnella diversifolia, 239
Neosharpiella aztecorum, 630
Nephroma arcticum, 405; cellulosum, 110; expallidum, 405; microphyllum, 108; parile, 405
New Hampshire, Sphagnum angermanicum in, 616
New Mexico, lichens in relation to the Four Corners power plant, 20
NICHOLS, G. E. (portrait), 511*
Nigeria, mosses from, II, 76
Nitrogen-fixing lichens, impact of emissions from coal-fired electricity generating facilities on, 54
nordivaricatic acid, 482
norgangaleoidin, 292+
NORRIS, D. H. & H. ROBINSON. The systematic position of Bryobrothera crenulata, 305
norstictic acid, 482
Northwest Territories, bryophytes new to, 409
Notas briologicas (Musci) IV y V, 99
Notholigotrichum mexicanum, 629; trichodon, 245
Notothylas anaporatus, 626, 637; dissecta, 625+, 627*; indica, 118, 146; levieri, 626, 628, 637; pandei, 628, 637; pfleidereri, 628
Nowellia curvifolia, 29+
Nucleosides and cyclic nucleotides, antagonism of cytokinin induced callus in Pylaisiella selwynii by, 47
nutritional conditions of sexual reproduction in Riccia, effect of, 37

Ocellularia americana, 125, 135; astroidea, 108; barroensis, 108; bullata, 108; congesta, 108; dactylifera, 108; dactyliza, 108; diacida, 108; dissuta, 108; domingensis, 134; esslingeri, 108; karnatakensis, 108; landronii, 108; lathraea, 135; minuta, 108; minutula, 108; neei, 108; pachystoma, 108; patwardlanii, 108; protocetrarica, 108; pseudoschizostoma, 108; pulverulenta, 108; referta, 108; subtilis, 125, 135; subwrightii, 108; unisepta, 108; verrucosa, 108; viridis, 108
Ochrobryum obtusifolium, 225
Ochrolechea androgyna, 405; frigida, 405; geminipara, 405; gotatodes, 405; grimmiae, 405; inaequalatula, 405; parella, 502; upsaliensis, 405
Octoblepharum albidum, 225, 439, 444, 479, 495; cocuiense, 216, 225; pellucidum, 225; pulvinatum, 225, 479; stramineum, 216
Odontoschisma elongata, 411; macounii, 162+; prostratum, 300, 484, 609, 610
Oedipodium griffithianum, 99+, 100*, 409+, 517
O'FLAHERTY, L. M. & G. W. FREEMAN. Eleven mosses new to Illinois, 609
Ohio, cold injury to lichens in southern, 620
OLARINMOYE, S. O. & A. EGUNYOMI. Mosses from Nigeria II. Collections from the Obudu cattle ranch, 76

Oligotrichum erosum, 245; falcatum, 409, 416; hercynicum, 410
Omphalanthus wallisii, 633
Omphalodiscus virginis, 23
Omphalodium arizonicum, 23
Oncophorus virens, 516
Opegrapha diaphora, 473; filicina, 88; lambinonii, 110
β-orcinol depsides, 154
Oreas martiana, 162, 170, 410
Oregon, new mosses from Crater Lake National Park, 82
Oreoweisia bogotensis, 224; laxifolia, 588; ligularis, 224, 253; serrulata, 516
Oropogon loxensis, 197, 315
Orthodicranum flagellare, 224, 439
Orthodontium confine, 229, 253; lineare, 515; longisetum, 229; osculatianum, 216; pellucens, 229, 253
rthorrhynchum chilense, 524; elegans, 519
Orthostichidium guayanense, 235, 252, 253; pentagonum, 253, 436
Orthostichopsis auricosta, 234; tetragona, 440, 480
Orthothecium intricatum, 414
Orthotrichaceae of Mexico, new taxa and new combinations in, 1
Orthotrichum affine, 264; alpestre, 264; anomalum, 29, 36, 519, 540, 549, 557, 611; apiculatum, 14; aristatum, 233, var. minus, 233; bartramii, 4; bolanderi, 264; consimile, 264; elongatum, 233; exiguum, 18; flowersii, 264, 266, 495; **hortonae**, 1+, 3*, 17*; laevigatum, 264; longirostre, 253; lozanoi, 4, var. lutescens, 4; lyellii, 264; malacophyllum, 4; mandonii, 233; mucronifolium, 14; obtusifolium, 275; ohioense, 486; pellucidum, 496; penicillatum, 233; pumilum, 275, 290, 611; punctatum, 6, 7; pycnophyllum, 4, 233, 253; pylaisii, 264; recurvans, 4, 233, 253; rufescens, 233, 253; rupestre, 142, 152, 264; sharpii, 3*, 4+, 17*; speciosum, 264; stellatum, 275, 609, 611; strangulatum, 486, 611; tenellum, 264; tuberculatum, 233; wallisii, 233
Osculatia columbica, 216, 229, 247, 253
OSORIO, H. S. Contribution to the lichen flora of Uruguay XI. Lichens from Arazatí, San José Department, 472
Oxyrrhynchum clinocarpum, 253; praelongum, 241, 250; remotifolium, 241, 250, 253; rugisetum, 241
Oxystegus cylindricus, 145, 147, 151, 439; tenuirostris, 499, 550, 554, var. gemmiparus, 499
Ozark National Forest, bryophytes of, 483
ozone fumigations, gross photosynthetic response of lichens to, 280

Pachyospora verrucosa, 405
Pallavicinia lyellii, 144, 300, 484
Paludella squarrosa, 412
Pannaria pityrea, 396+
Pannoparmelia wilsonii, 316
PAOLILLO, D. J., JR. On the lipids of the sperm masses of three mosses, 93
Papillaria appressa, 253; deppei, 235, 436; fusco-viridis, 235, 253; imponderosa, 235, 251; nigrescens, 235, 252, 253, 439, 480; penicillata, 235; subsquamatula, 235
Paraleucobryum densifolium, 223, 253, var. congestum, 223; var. latilimbatum, 223; longifolium, 275, 412
parietin, 302
Parmelia almquistii, 396+; alpicola, 405; arseneana, 23, 27; borrerioides, 473; caperata, 174, 502, 620, 621; centrifuga, 405; chlorochroa, 23; conspersa, 23, 174; crinita, 620; cryptochlorophaea, 135; cumberlandia, 23, 501; dierythra, 23; disjuncta, 405; elegantula, 23; exasperatula, 405; fraudans, 396+; fuliginosa, 111; glabratula subsp. fuliginosa, 106; goebelii, 135; hypoleucites, 23; hypotropa, 620; imatatrix, 174; incurva, 405; infumata, 23, 405; kurakawa, 23; laevigatula, 135; lineola, 23; livida, 620; margaritata, 620; mexicana, 23, 27; michauxiana, 620; microsticta, 473; omphalodes, 405; panniformis, 405; physodes, 505, 574; plittii, 23; praesignis, 564; protomatrae, 161; psoromifera, 23; pulla, 159; reticulata, 135, 620, 621; rudecta, 620, 621; saxatilis, 405; separata, 405; septentrionalis, 405; signifera, 174+; stuppea, 620; stygia, 405; subaurifera, 23; subcrinita, 135, 620; subdecipiens, 23; subisidiosa, 620, 621; subolivacea, 23; subrudecta, 620; substygia, 23, 24, 405; sulcata, 107, 111, 175, 280+, 315, 316, 405, 564+; tasmanica, 175; texana, 135, 620; tiliacea, 314; ulophyllodes, 23; ultralucens, 135
Parmeliopsis aleurites, 136; ambigua, 136; halei, 126, 136; hyperopta, 396+; pallescens, 136; subambigua, 125, 136
Parmotrema cetratum, 473; eciliatum, 473; subsumptum, 473; ultralucens, 135
Pellia epiphylla, 144, 145, 582, 636, var. borealis, 118
Peltigera aphthosa, 319, 405; canina, 23, 180, 316, 405, var. praetextata, 574; lepidophora, 405; membranacea, 406; polydactyla, 180, 406, 574; praetextata, 406; rufescens, 406; spuria, 406; venosa, 406
Peltula bolanderi, 23; obscurans var. hassei, 23
penicillin-resistant bacteria, 141
perlatolic acid, 195
Peromnium daguense, 228, 253; jamesonii, 228, 253
peroxyacetylnitrate fumigation, response of lichens to, 564
Pertusaria amara, 125, 136; bryontha, 406; dac-

tylina, 406; multipuncta, 136; panyurga, 406; pulchella, 473; saximontana, 23; subplicans, 406; tuberculifera, 137; velata, 137; xanthodes, 137
Petalophyllum ralfsii, 300
PETERSON, P. J. *See* M. A. S. BURTON & P. J. PETERSON, 594
Phaeoceros fimbriatus, 633; laevis, 300, 484, 634; miyakeanus, 634; qualaquizana, 633
Phaeographina arechavalatae, 473; atrovermicularis, 320; coriaria, 320; galactinum, 109; quassiaecola, 125, 137
Phaeographis lyellii, 125, 137; mordenii, 320; pezizoidea, 473
Phaeophyscia adiastola, 107; cernohorski, 107; chlorantha, 109; decolor, 107; endococcinodes, 107; erythrocardia, 107; hirsuta, 107, 109; hirtuosa, 107; hispidula, 107, 109; imbricata, 107; insignis, 109; laciniata, 315; leana, 107; pusilloides, 107; rubropulchra, 107, 109, 125, 138; trichophora, 107
Phaeorhiza nimbosa, 504; sareptana, 504, var. sphaerocarpa, 504
Phaeotrema psoromicum, 108; scabridum, 108; vulcani, 108
Phascum curvicolle, 542; cuspidatum, 264, 540, 542, 557, 609; floerkeanum, 542; hyalinotrichum, 265
Phenology of selected bryophytes in southern Illinois, 29
Philibertiella ditrichoidea, 516
Philonotis andina, 231; angulata, 231; crassinervia, 231; curvata, 231; elegantula, 231; falcata, 146; fontanella, 231; fontana, 74, 83, 265, 275, 486, 515, 622; glaucescens, 253, 436, 480; marchica, 232, 253, 486, 495, 611, 612; osculatianum, 216, 232; revoluta, 232; rigida, 498; scabrifolia, 232, 246; sphaericarpa, 232; striatula, 232; tenella, 253; umbonata, 232, 253; uncata var. gracilenta, 246; uncinata, 232, var. glaucescens, 232, 246, 247, 253, var. gracilenta, 232, var. gracillima, 232
Phoma truncata, 502
photochemical activities in liverworts, 286+
photoperiodism in bryophytes, 29+
photosynthesis in liverworts, 286+
photosynthetic assimilation rates in lichens, 171+
photosynthetic response of lichens to ozone, 280+
photosynthetic response of lichens to peroxyacetylnitrate fumigation, 564
Phyllodrepanium falcifolium, 230
Phyllogonium fulgens, 236, 437, 480; viscosum, 236
Phyllopsora corallina, 137
Physica aipolia, 23, 406; alba, 23, 473; adscendens, 406; biziana, 23; caesia, 23, 406; callosa, 23; carassensis var. obsessa, 473; cascadensis, 23; ciliata, 23; dubia, 23; endococcinea, 23, 406; halei, 23; hispidula, 23; intermedia, 23; orbicularis, 23; phaea, 24, 404; stellaris, 24; wainioi, 24
Physconia grisea, 24, 406; muscigena, 24, 406; pulverulacea, 316; pulverulenta, 316
Physcomitrella patens, 609
Physcomitrium immersum, 611; pyriforme, 275, 486, 492, 609, 611; sphaericum, 517
physiological responses in lichens, 171+
Pilophoron robustus, 396+
Pilopogon gracilis, 224, 437; laevis, 224, 253; longirostratus, 224; muelleri, 223; nanus, 224; paleaceus, 224; piliferus, 224, 253, var. proliferocomus, 224
Pilopogonella laevis, 253
Pilosium chlorophyllum, 242
Pilotrichella consimilis, 235, 253; flexilis, 235, 437; fusco-viridis, 253; longebarbata, 235, 253; nigricans, 253; pachygaster, 235, var. gracilis, 235; pentasticha, 235; remotifolia, 253; rigida, 235, 435, 480; tenuissima, 253; viridis, 235
Pilotrichum andersonii, 216; asperifolium, 236, 253; bipinnatum, 253; fendleri, 253; lindigii, 236, 253; lycopodium, 234, 253; macranthum, 235, 253; nitidum, 237, 253; patens, 253; ramosissimum, 237, 253, 480; sticticola, 235, 253; wallisii, 237, 253
Pinnatella minuta, 437, 444
Pireella cymbifolia, 480; mariae, 435; pohlii, 437, 441
Placopsis gelida, 406
Placynthium aspratile, 406
Plagiobryum zierii, 489
Plagiochasma appendiculatum, 145, 149, 150, 151, 152; intermedium, 632; microcephalum, 113; ruprestre, 300, var. volkii, 113
Plagiochila acanthophylla subsp. japonica, 484; artica, 162+; asplenioides, 114, 116, 166, 274, 484, var. subarctica, 165+; hattoriana, 310; ludoviciana, 484; major, 116, 634; porelloides, 116, 165+; undata, 484; yokogurensis, 112
Plagiomnium medium, 73; rhynchophorum, 230, 252
Plagiothecium andinum, 242; cavifolium, 275, 412, 486, 611; denticulatum, 142, 242, 519, 611; laetum, 275; lucidulum, 242; novogranatense, 242; planissimum, 437; pseudotenerum, 242, 253; roeseanum, 242; tenerum, 253; undulatum, 150, 494
planaic acid, 189+
Platydictya confervoides, 275, 290, 486, 611; sprucei, 514; subtile, 275, 511
Platygrapha leucophthalma, 89
Platygyriella helicodontoides, 434
Platygyrium repens, 275, 486, 498, 611
Platyhypnidium aquaticum, 241; riparioides, 241; subrusciforme, 435

Platyneuron laticostatum, 516
Pleospora peripherica, 502
Pleuridium bolanderi, 265; californicum, 265; nervosum, 598, 599; subulatum, 73, 611
Pleurochaete luteola, 227, 437; squarrosa, 486, 545
Pleuropus bonplandii, 241, 253
Pleurorthotrichum chilense, 519
Pleurotheliopsis erigens, 503
Pleuroweisia schliephackei, 545
Pleurozium schreberi, 242, 328, 343, 367, 599+
Poecilophyllum procumbens, 224, 253
Pogonatum abbreviatum, 245; aloides, 142; alpiniforme, 245, 253; alpinum, 275; andinum, 245; arcuatum, 245; brachyphyllum, 486, 611; cucullatum, 245, 253; cuspidatum, 245; microstomum, 143, 144; neglectum, 245; oligodus, 245, 253, 254; pensilvanicum, 438, 611; psilopiloides, 245; purpurascens, 245; robustum, 437; semipellucidum, 245, 253; subbifarium, 245, 253; subcontortum, 245; tortile, 245, 439; varians, 245; viride, 245, 253
Pohlia annotina, 265, var. decipiens, 275; bulbifera, 265, 414; cruegeri, 253; cruda, 81, 265, 275; crudoides, 411; elongata, 265, 412, 515; flexuosa, 229; fusifera, 229; integridens, 229, 247, 256, 435, 529; leptodontium, 230; lutescens, 478; nutans, 99, 265, 364, 486, 611; papillosa, 230; paucifolia, 230; pulchella, 478; salaminae, 230; subcurvata, 230; wahlenbergii, 265, 275, 486, 609, 611
Polyblastia gothica, 406; hyperborea, 406; melanospora, 502; scotinospora, 502; sendtneri, 406; sprucei, 396; theolodes, 406
Polyblastiopsis swinscowii, 505
Polychidium dendriscum, 125, 138
Polysaccharides in Usnea cavernosa, ultrastructural localization of, 533
Polysporina cyclocarpa, 506; dubia, 506; simplex, 506
Polytrichadelphus abriaquiae, 245; aristatus, 245; ciliatus, 245; ericoides, 253; giganteus, 245; longisetus, 245, 247, 253, 254; purpureus, 245; rubiginosus, 245; trianae, 245, 253
Polytrichastrum tenellum, 245, 253
Polytrichum aloides, 366, 519; alpestre, 325+; alpinum, 366; andinum, 245, 253; antarcticum, 519; antillarum, 245, subsp. substrictum, 245; aristiflorum, 245; attenuatum, 93; australe, 519; ciliatum, 245; commune, 141, 151, 245, 358, 439, 486, 519, 611; convolutum, 245; cucullatum, 245; ericoides, 245; equisetiforme, 245; formosum, 275; giganteum, 245; hercynicum, 519; jamesonii, 245, 254; juniperinum, 73, 141, 245, 254, 258, 265, 438, 486, 611; longisetum, 245, 254, 414; magellanicum, 519; neglectum, 245, 254; ohioense, 275, 486, 611; piliferum, 74, 265, 366; psilopiloides, 245, 254; purpurascens, 245, 254; rhynchomitrium, 245, 254; semipellucidum, 254; subcontortum, 245, 254; thysanomitrium, 245, 254; trianae, 245, 254; trichodon, 245, 254; varians, 245, 254
population biology of Polytrichum alpestre, 325+
Porella acutifolia subsp. tosana, 310; campylophylla subsp. lancistipula, 310; chinensis var. decurrens, 310; obtusata forma macroloba, 311; pinnata, 300, 484; platyphylla, 11, 115, 484, 581, 610; platyphylloidea, 115, 274, 300, 484
Porina thaxteri, 88
Porothamnium elatulum, 236; expansum, 236; fasciculatum, 236; flagelliferum, 236; lancifrons, 236; neckeraeforme, 236; sparsiflorum, 236, 251; substriatum, 236
Porotrichodendron mahahaianum, 518; nitidum, 237; robustum, 237; superbum, 237, 254
Porotrichum cobanense, 256, 437; decompositum, 254; fasciculatum, 254; flagelliferum, 254; lancifrons, 254; lindigii, 236; longirostre, 236, 437, 518; minus, 237; mutabile, 237, 254; neckeraeforme, 254; nitidum, 254; plicatulum, 237, 480; scoposum, 237; sparsiflorum, 254; stolonaceum, 237, 254; substriatum, 254; superbum, 237, 254; variabile, 254
Potamium pulchellum, 254
Pottia arizonica, 265, 542, var. mucronulata, 265; bryoides, 265, 542; cucullata, 227; davalliana, 542, 555, subsp. commutata, 542; fosbergii, 265, 266, 542; heimii, 542, 549; intermedia, 542; lanceolata, 454, 542; mutica, 542; nevadensis, 542; recta, 542; starkeana, 265, 542+; truncata, 290, 520, 542; viridifolia, 542; wilsonii, 542, subsp. asperula, 542, var. crinita, 542
Preissia quadrata, 484
Prionodon densus, 234, 253, 254, 439; dichotomus, 234, 257, 437; divaricatus, 234, 254; flagellaris, 234, 254; fusco-lutescens, 234, 254; laeviusculus, 254; luteo-virens, 234, 254, 257; lycopodioides, 234; muelleri, 234, 254; pinnatus, 234; undulatus, 234, 254
PRIOR, P. V. *See* D. E. JACOBSON & P. V. PRIOR, 622
propaguliferous Ceratodon purpureus, 474
Protoblastenia rupestris, 406
Pseudephebe minuscula, 406; pubescens, 406
Pseudobraunia californica, 265
Pseudobryum cinclidioides, 414
Pseudocryphaea flagellifera, 437
Pseudocyphellaria aurata, 138; australiensis, 174+; billardierii, 174; carpaloma, 174; colensoi, 174; crocata, 174+; delisei, 171+; fav-

eolata, 174; flotowiana, 174; glabra, 172; granulata, 505; hirta, 174; mougeotiana, 174; physciospora, 505; rubella, 174
Pseudoevernia furfuracea, 500
Pseudodistichium austro-georgicum, 516
Pseudoleskea patens, 265; radicosa, 265
Pseudoparmelia crozalsiana, 473; cryptochlorophaea, 135; exornata, 318; texana, 135
Pseudosymblepharis angustata, 545; mauiensis, 80; schimperiana, 437
Psilopilum mexicanum, 629, 631; trichodon, 254
Psoroma hypnorum, 406; leprolomum, 315
Pterigynandrum consanguineum, 254; cylindricaule, 242, 254; longisetum, 242, 254; squarrosum, 254
Pterobryum angustifolium, 480; densum, 437; duplicatum, 254; lindigii, 254
Pterobryopsis mexicana, 435
Pterogonidium pulchellum, 79, 243
Pterogonium fulgens, 254; gracile, 265; pulchellum, 243, 254
Pterygoneurum lamellatum, 542; ovatum, 542; subsessile, 542
Ptylidium ciliare, 163; pulcherrimum, 167
Ptychomitrium crispatum, 454; incurvum, 486; lindigii, 232; lobuliferum, 232; polyphyllum, 517; serratum, 435, 441
Ptychomnion aciculare, 520
Puerto Rico, chromosome number of Callicostella depressa, 103+
PURSELL, R. A. On Fissidens. Nomenclatural and distributional notes on some American species, 58
PURSELL, R. A. & E. HEGEWALD. On Fissidens in Peru, 267
Pycnolejeunea contigua, 634
Pylaisia polyantha, 557
Pylaisiella falcata, 244; polyantha, 539; selwynii, 47+, 51*, 275, 486
Pyrenotrichum splitgerberi, 91, 92, 125, 138
Pyrenula clandestina, 505

Racomitrium conterminum, 227, 254; crispipilum, 227, 254; crispulum, 249, 254; cucullatifolium, 228; dichelymoides, 228; lanuginosum, 30, 36, 71, 228, 366, 367, 539, 558
Radula boryana, 119; caduca, 637; campanigera subsp. obiensis, 637; complanata, 310; inouei, 637; iwatsukii, 637; lindenbergiana, 114; mollis, 484; obconica, 484; ocellata, 637; philippinensis, 637; prolifera, 162+; retroflexa var. fauciloba, 637; van-zantenii, 637; variabilis, 112, 310; verrucosa, 637; yangii, 637
Ramalina almquistii, 406; americana, 502; anceps var. peranceps, 319; celastri, 473; complanata, 473, 502; duriaei, 505; farinacea, 106; fastigiata, 502; maciformis, 65, 71, 179, 503, 574; muciformis, 515; protecta, 314; sharpii, 319; sinensis, 24; stenospora, 502; usnea, 319
RAO, K. R., N. R. KUMAR & A. N. REDDY. Studies of photosynthesis in some liverworts, 286
Rauiella praelonga, 435; subcatenulata, 435
RAVENEL, H. W. (portrait), 511*
Recent literature on hepatics and anthocerotes—83, 112; —84, 310; —85, 632
Recent literature on lichens—101, 105; —102, 314; —103, 500
Recent literature on mosses—13, 494
Reboulia hemisphaerica, 145, 149, 151, 286+, 300, 484, 610
REDDY, A. N. *See* K. R. RAO, N. R. KUMAR & A. N. REDDY, 286
REDFEARN, P. L. Bryophytes of the Interior Highlands of North America XVI. Species of the Ozark National Forest, Sycamore Division, and adjacent areas in Stone County, Arkansas, 483
REESE, W. D. Calymperes (Musci) from western Amazônia: Brazil and Bolivia, 559
Renauldia cochlearifolia, 480; hildebrantielloides, 520; mexicana, 435
respiration in Fontinalis duriaei, 382
Rhabdoweisia crenulata, 224, 259, 583, 584, 585; crispata, 81, 274, 411, 412, 585; cyathicarpa, 254; denticulata, 585; fugax, 224, 254, 256, 630; lindigiana, 224, 254
Rhachithecium perpusillum, 499, 510
Rhacocarpus humboldtii, 234, 527, var. orbiculatus, 234; purpurascens, 234, 249
Rhacopilum intermedium, 233; mnioides, 520; tomentosum, 233, 439
Rhamphidium macrostegium, 520
Rhaphidorrhynchium aurescens, 520; decumbens, 243; erythropodium, 243; fragilirostrum, 243; lindigii, 243, 435; roridum, 243; schlimii, 243; sticticola, 243; subscabrum, 243
Rhaphidostegium galipense, 254; loxense, 243, 254; subloxense, 243, 254; subsimplex, 254
Rhegmatodon filiformis, 435
Rhizocarpon alpicola, 503; chioneum, 406; crystallinum, 396; crystalligenum, 406; disporum, 24, 406; eupetraeoides, 406; eupetraeum, 406; geographicum, 109, 406, 503, subsp. arcticum, 406; grande, 406; hochstetteri, 396+; inarense, 396+; intermedium, 396+; obscuratum, 502; polycarpum, 396+; pusillum, 406; riparium, 24; rittokense, 406
Rhizogonium lindigii, 230; mnioides, 230, 252; spiniforme, 230, 439, 480
Rhizomnium andrewsianum, 410; magnifolium, 495, 497; pseudopunctatum, 74
Rhizoplaca chrysoleuca, 24, 406; glaucophana, 506; haydenii, 506; marginalis, 506
Rhodobryum beyrichianum, 230, 435, 480; gi-

ganteum, 143, 146; grandifolium, 230; huillense, 498; mniopsis, 230; ontariense, 74, 412, 416; pycnopyxis, 230; roseum, 412, 416, 439, 486, 515, 612
Rhynchostegiella attenuata, 241; compacta, 265; tenella, 515
Rhynchostegiopsis flexuosa, 437
Rhynchostegium aquaticum, 254; chrysophylloides, 241; conchophyllum, 241; confertum, 515; deplanatiforme, 242, 254; pulchellum, 254, 375, 486; remotifolium, 254; riparioides, 254; scariosum, 241, 435; semiscabrum, 254, 439; serrulatum, 275, 486, 492, 612; subrotundum, 241; tocaremae, 241; ulicon, 241
Rhytidiadelphus loreus, 150; novogranatensis, 244; squarrosus, 412, 496, 596, 597; triquetrus, 74
Riccardia aberrans, 633; amazonica, 633; bogotensis, 633; cervicornis, 633; chamaedryfolia forma major, 118; ciliolata, 633; columbica, 633; latifrons, 300; lepidomitra, 635; metzgeriaeformis, 633; multifida, 484; papillata, 633; pinguis, 582; squarrosa, 633; wallisii, 633
Riccia abuensis, 37; arnellii, 98; californica, 636; campbelliana, 636; canaliculata, 37+; curtisii, 98; dictyospora, 300; discolor, 149; duplex, 37+, 117; fluitans, 37+, 117, 300, 609, 610, 634; frostii, 45, 46; glauca, 609, 610, 635; ianthina, 311; liaoningensis, 633; limnicola, 311; macallisteri, 300; media, 37; pseudofluitans, 633; rhenana, 37+; sorocarpa, 634, 636; stricta, 37; trichocarpa, 636
Ricciocarpus natans, 114, 274, 300, 610, 636
Riella affinis, 635, 636; americana, 118; helicophylla, 636
Rigodium gracile, 435; implexum, 515; solutum, 237; toxarioides, 237; toxarion, 237
RINNE, R. J. K. *See* P. BARCLAY-ESTRUP & R. J. K. RINNE, 599
Rinodina archaea, 24; bischoffii, 24; calcarea, 24; colobinoides, 125, 138; exigua, 24; granuligera, 125, 138; hyperborea, 406; iowensis, 138; laevigata, 396+; milliaria, 125, 138; nimbosa, 504; occidentalis, 406; ochrocea, 125, 138; sareptana, 504; septentrionalis, 502; turfacea, 406
ROBINSON, H. *See* D. H. NORRIS & H. ROBINSON, 305
RUNDEL, P. W., G. C. BRATT & O. L. LANGE. Habitat ecology and physiological response of Sticta filix and Pseudocyphellaria delisei from Tasmania, 171
RUSSELL, J. L. (portrait), 511*

Saelania glaucescens, 73, 456
SANFORD, G. R. Temperature related growth patterns in Amblystegium riparium, 525
Sarcographa tricosa, 138
Sarcogyne clavus, 125, 138; privigna, 24; regularis, 406; simplex, 138
Sarconeurum glaciale, 84, 87; tortelloides, 84+, 478
Sauloma tenella, 517
Scapania americana, 181+; apiculata, 182+; bolanderi, 181+; brevicaulis, 182+; carinthiaca, 183+; compacta, 181; crassiretis, 184+, 411, 637; curta, 181+; cuspiduligera, 182+; degenii, 118, 183+, var. dubia, 184; glaucocephala, 181+; granulifera, 181; gymnostomophila, 182+; helvetica, 182; hians, 186, subsp. salishensis, 184+; hyperborea, 183+; irrigua, 181, subsp. rufescens, 184; jensenii, 118; massalongoi, 184; mucronata, 182+, var. praetervisa, 183; nemorea, 184+; nemorosa, 181+, 274, 301, 484, 610; oakesii, 181+; obcordata, 183+; paludicola, 182+; paludosa, 182+, 411; parvifolia, 184; praetervisa, 184; saxicola, 183+; scandica, 182+, 411; simmonsii, 162+, 183+, 637; spitzbergensis, 162+, 182+; subalpina, 119, 181+, 411; uliginosa, 168, 181+; umbrosa, 181+; undulata, 114, 181, 484, 594+, var. oakesii, 183+
SCHERTLER, M. M. Development of the archegonium and embryo in Lophocolea heterophylla, 756
Schimperobryum splendidissimum, 517
Schistidium alpicola, 73, 74
Schistochila philippinensis, 119
Schistomitrium obtusifolium, 254
Schistostega pennata, 274, 495
Schliephackea meteorioides, 224; prostrata, 224
Schlotheimia brachyrhyncha, 6; jamesonii, 437; rugifolia, 438, 492
SCHNEIDER, A. (portrait), 512*
SCHOFIELD, W. B. *See* J. D. GODFREY & W. B. SCHOFIELD
SCHWENDENER, S. (portrait), 124*
Schwetschkeopsis elongata, 494; fabronia, 486
Sciaromium compissatum, 514; crassinervatum, 240
Scleropodium caespitans, 265; californicum, 265; julaceum, 265; obtusifolium, 265; tourettei, 265
Scouleria aquatica, 102, 517; marginata, 100*, 101, 102; patagonica, 99+, 100*
Scoliciosporum chlorococcum, 506; pruinosum, 506; sarcothamni, 506; schadeanum, 506
Seligeria calcarea, 415; lindigiana, 224, 254; oelandica, 412; subimmersa, 274, 410
SELKIRK, P. M. Effect of nutritional conditions on sexual reproduction in Riccia, 37
Sematophyllum adnatum, 609, 612; agnatum, 243, 254; aureosulfureum, 243; caespitosum, 81, 243, 250, 254, 255, 439; chrysostegum, 243; crassiusculum, 243; crinitifol-

ium, 254; crithmifolium, 243, 254; cucullatifolium, 243, 254; cuspidiferum, 437; decumbens, 243, 254; demissum, 286, 520; erythropodium, 243, 255; flavidum, 243; fluviale, 243, 255; fragilirostrum, 255; galipense, 243; hawaiiense, 81; insularum, 437; leptothecium, 243, 255; lindigii, 255; loxense, 243, 255; pennellii, 243; pungens, 255; roridum, 243, 255; sericifolium, 437; subbrachycarpum, 243; subsimplex, 243; subscabrum, 255; tequendamense, 243, 250, 254
Semibarbula orientalis, 144
SEN, S. P. *See* R. D. BANERJEE & S. P. SEN
SÉRUSIAUX, E. Foliicolous lichens from Southeastern United States, 88
sexual reproduction in Riccia, effect of nutritional conditions in, 37
SEYER, S. C. New mosses from Crater Lake National Park, Oregon, 82
SHERIDAN, R. P. Impact of emissions from coal-fired electricity generating facilities on N_2-fixing lichens, 54
SHOWMAN, R. E. Cold injury to lichens in southern Ohio, 620
SIGAL, L. L. *See* T. H. NASH III & L. L. SIGAL, 280
SIGAL, L. L. & O. C. TAYLOR. Preliminary studies of the gross photosynthetic response of lichens to peroxyacetylnitrate fumigation, 564
SINGH, D. K. *See* R. UDAR & D. K. SINGH, 625
Skottsbergia paradoxa, 516
Solenostoma crenulatum, 594+
Solorina crocea, 316, 406; saccata, 406; spongiosa, 406
Sonoran desert, desiccation of mosses in, 65
South Dakota, additions to the mosses, 72
sperm masses of three mosses, the lipids of, 93
Sphaerocarpus donnellii, 44, 46; texanus, 609, 610
Sphaerophorus fragilis, 406; globosus, 398+; melanocarpus, 174; tener, 174
Sphaerothecium comosum, 225, 255, var. elatius, 225; phascoides, 225, 255
Sphagnum acutifolium, 219, 255; andinum, 219, 255; angermannicum, 616+; angustifolium, 297, 412; annulatum, 295+, var. porosum, 295; apollinairei, 220, 255; balticum, 297; bartlettianum, 198+, 199*, 297; capillaceum, 219, 255, 256, 275; capillifolium, 219, 255, 609, 610, 612, var. tenerum, 609, 610, 612; compactum, 219, 297, 486; contortum, 410; coryphaeum, 219, 255; cuspidatum, 151, 219, 255, 297; cyclophyllum, 219, 617; cymbifolium, 220, 255; erythrocalyx, 219; fallax, 297; fitzgeraldii, 297; flavicomans, 203, 297; fuscum, 297, 328+, 497; girgensohnii, 297; henryense, 297; imbricatum, 297, 492; jensenii, 295+, 296*; lehmannii, 255, var. equiporosum, 219; lenense, 297; lescurii, 297; limbatum, 219, 255; lindbergii, 297, 617; macrophyllum, 297; magellanicum, 153, 219, 255, 297, 328, 339, 492, 498, 616; majus, 295+, 616; medium, 219, 255, var. albescens, 219, var. pallescens, 219, var. purpurescens, 219; mendocinum, 298; meridense, 219, 255; nemoreum, 203, 255, 297; obtusum, 298; oxyphyllum, 220; palustre, 151, 220, 255, 297, 492, 521; papillosum, 297, 616; perichaetiale, 201, 297, 491; platyphyllum, 412; plumulosum, 203; portoricense, 142, 143, 152, 297; pulchricoma, 220, 255; pulchrum, 297; pylaesii, 220, 297, 617; quinquefarium, 198+; recurvum, 201+, 220, 255, 297, 328; riparium, 297; rubellum, 198+, 297, 366; russowii, 616; sancto-josephense, 220; sparsum, 220, 255; squarrosum, 74, 297, 412; strictum, 142, 201, 297; subfulvum, 203, 297; subsecundum, 220; subtile, 198+, 297; tenellum, 617; tenerum, 201, 297; teres, 297; torreyanum, 297; trinitense, 297; wallisii, 219, 255; warnstorfianum, 198; warnstorfii, 74, 198+
Sphinctrina tubaeformis, 125, 138
SPIESS, L. D. Antagonism of cytokinin induced callus in Pylaisiella selwynii by nucleosides and cyclic nucleotides, 47
Splachnobryum obtusum, 228, 520
Splachnum ampullaceum var. brevisetus, 496; pensylvanicum, 495; turbinatum, 496; weberbaueri, 228
Sporastatia testudinea, 406
Sporledera lindigiana, 221, 255; subenervis, 221, 255
Squamarina lentigera, 24
squamatic acid, 154
Squamidium filiferum, 235; leucotrichum, 235, 252, 255, 437; nigricans, 235, 253, 437, 480; ternstroemiae, 235
Staurothele catalepta, 24; clopima, 24; fissa, 396+; rufa, 24
STEERE, W. C. Some mosses from British Honduras, 478
STEERE, W. C. The chromosome number of Callicostella depressa, 103
Steereella linearis, 613+, 614*
Stegonia latifolia, 543
Stenodesmus latifolius, 239
Stereocaulon alpinum, 397+; glareosum, 406; meyeri, 110, subsp. atlanticum, 110; paschale, 106, 108, 316, 406; ramulosum, 110, 174+, 189+; tomentosum, 407
Stereodon affinis, 255; amabilis, 255; hamatus, 244, 255; spiripes, 255
Stereophyllum leucostegium, 481; radiculosum, 437, 480
Sticta dufourei, 505; filix, 171+; fuliginosa, 174; limbata, 174; weigelii, 396+

Stictolejeunea iwatsuki, 312
Stirtonia alba, 125, 139
Stokesiella praelonga, 255, 265, 515
STONEBURNER, A. & R. WYATT. Three Big Thicket bryophytes new to Texas, 491
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—83, 112; —84, 310; —85, 632
STRASBURGER, E. A. (portrait), 124*
Streptopodon calymperes, 227, 247, 255; cavifolius, 227, 435; complanata, 88; elegans, 88, 91; erythrodontus, 227; feei, 91; latifolius, 227, 255; lindigii, 227, 255; matudianus, 434; rigidus, 227, 255; setiferus, 227, 255
sulfur dioxide, effect on lichens, 20+, 54+
SULLIVANT, W. S. (portrait), 512*
Symblepharis lindigii, 225
Symphyogyna aspera, 582
Syrrhopodon crispatus, 226, 255; cryptocarpos, 225, 559+; elongatus var. glaziovii, 497; flammeo-nervis subsp. annamensis, 498, subsp. lilliei, 498; fragilis, 227, 255; gaudichaudii, 225, 255, 437; helicophyllus, 225, 629; hornschuchii, 225, 260, 561; incompletus, 225, 438, 479, 559, 561; inflexus, 225, 255; iridans, 225, 255; leprieurii, 225, 255, 561; ligulatus, 562; lycopodioides, 225, 437; parasiticus, 225, 493, 559, 562; prolifer, 225, 437, 562, var. acanthoneuros, 562, var. cincinnatus, 497, var. papillosus, 497, 562, var. tenuifolius, 497; pusillus, 225, 255; quintasii, 225, 258, 439; rigidus, 225; rupestris, 255; simmondsii, 225, 560, 562; spiculosus, 498; texanus, 492; xanthophyllus, 560, 563
Systegium lindigii, 224, 255

Takakia ceratophylla, 115, 312
TAN, B. C. (review), P-C. C. CHEN et al., Genera Muscorum Sinicorum, 638
Tapellaria epiphylla, 88+, 89*
Targionia hypophylla, 117, 149, 151, 298+
Tasmania, habitat ecology and physiological response of Sticta filix and Pseudocyphellaria delisei from, 171
Taxiphyllum deplanatum, 486, 612; taxirameum, 290, 291, 486
Taxithelium nepalense, 147; planum, 243, 481
TAYLOR, O. C. *See* L. L. SIGAL & O. C. TAYLOR, 564
Tayloria altorum, 228; henryae, 496; jamesonii, 255; mayorii, 228, 255; moritziana, 255; scabriseta, 228; splachnoides, 521
Telaranea nematodes, 491
Teloschistes capensis, 270+*; chrysophthalmus, 473; flavicans, 305; villosus, 505
temperature effects on assimilation and respiration in the Fontinalis duriaei-periphyton association, 382
temperature related growth patterns in Amblystegium riparium, 525
Tetralophozia setiformis, 118
Tetraphis pellucida, 275
Tetraplodon angustatus var. integerrimus, 496; balticus, 495; blyttii, 495; mnioides, 228, 430; urceolatus var. longisetus, 496
Tetrapterum australe, 520
Texas, notes on hepatics, 298; Big Thicket bryophytes new to, 491
Thamniopsis killipii, 239; pendula, 239
Thamnolia subuliformis, 407; vermicularis, 319, 398+
thamnolic acid, 482
Thelia asprella, 612, hirtella, 486; lescurii, 486, 492, 612
Thelidum acrotellum, 407
Thelopsis flaveola, 77, 78, 502; inordinata, 77, 78, 502; isiaca, 77; melathelia, 77, 78, 502; rubella, 77, 78, 502
Thelotrema acharium, 505; demersum, 505; exile, 108; hiatum, 108; maximum, 108, sanfordianum, 125, 139; wightii, 140
THOMSON, J. W. *See* T. J. MOSER, T. H. NASH III & J. W. THOMSON, 393
Thuidium antillarum, 240, 255, 437; arzobispoae, 240, 250; carantae, 240; ciliarifolium, 498; ciliatum, 240; complanum, 240; delicatulum, 240, 255, 275, 486, 612; frontinoae, 240; gratum, 240, 497; involvens, 240; minutulum, 240, 250, 439, 518; peruvianum, 240; pseudo-delicatulum, 240; pygmaeum, 291, 486; recognitum, 74, 240, 486, 612, var. delicatulum, 240, 250, 439, subsp. philibertii, 255; schistocalyx, 480; tamariscinum, 240; tomentosum, 435; urceolatum, 240, 255
Thysanomitrium muelleri, 223, 255; phascoides, 225, 255; richardii, 255
Tichothecium pygmaeum, 407
Timmia megapolitana, 486, 612; norvegica, 415
Timmiella anomala, 265, 439, 546, 555; barbuloides, 546; crassinervis, 265; flexiseta, 546
TIXIER, P. À propos de quelques mousses nouvelles pour le sud de Madagascar, 598
TIXIER, P. Contribution to the knowledge of the genus Cololejeunea VIII. Some new species of Malagasy Cololejeunea, 602
Tomenthypnum falcifolium, 412, 416; nitens, 366, 412
Toninia caeruleonigricans, 24, 407; candida, 24; cinerovirens, 24; lobulata, 407; tristis, 24, 407
Tortella caespitosa, 520; flavovirens, 545, var. viridiflava, 545; fragilis, 84+, 476, 545, 552, var. fragilis, 84+, var. tortelloides, 84+, 476, 478; humilis, 439, 486, 545, 609, 612; inclinata, 545; inflexa, 545; japonica, 545, 552;

mollissima, 437; tortelloides, 84+; tortuosa, 74, 79, 86, 275, 291, 545, 552
Tortula **ammonsiana**, 466+*; amplexa, 265, 543, 550; andicola, 227; angustifolia, 631; apiculata, 226, 255; arcuata, 226, 255; bartramii, 266, bistratosa, 266; bogotensis, 227, 255; bolanderi, 266; brevipes, 266; buyssonii, 543; californica, 266; canescens, 543; caroliniana, 437; cuneifolia, 543; decidua, 227, 255; decolorans, 255; densifolia, 630; ferganensis, 543; fragilis, 227, 246, 255, 266, 543, 550; glaucescens, 255; goudotii, 227; grandiretis, 495; guepinii, 543; handelii var. ferganensis, 496; herzogii, 631; inermis, 67+, 266, 543; intermedia, 543; kingii, 227; laevipila, 266, 469+, 543, 550, var. meridionalis, 470, var. propagulifera, 471, 472; wachteri, 471, 472; latifolia, 543; lingulata, 543; longirostris, 255; marginata, 543; mniadelphus, 630; mniifolia, 437; mucronifolia, 266, 543; muralis, 266, 291, 539, 543, 551; norvegica, 266, 543; novogranatensis, 255; obtusissima, 266; pagorum, 67+, 73, 469+, 470+, 486; papillosa, 266, 291; pinchinchensis, 227, 246; plinthobius, 543, 551; princeps, 266, 543; propagulosa, 499, 510; rectifolia, 543; revolutifolia, 543; revolvens, 543; ruralis, 58, 67+, 142, 266, 543, var. crinita, 266, var. hirsuta, 266, var. ruraliformis, 543; scotteri, 499; schlimii, 255; sinensis, 543; stenocarpa, 255; subulata, 266, 520, 543, var. angustata, 543; thianschanica, 543; transcaspica, 543; trianae, 227, 255; vahliana, 543; virescens, 543, 550
trace element accumulation in a feather moss and in soil near a Kraft paper mill in Ontario, 599
Trachypodopsis serrulata var. crispatula, 79
Trachypus bicolor, 81, var. viridulus, 439
Trapelia coarctata, 506
TRAVERSE, A. (review), S. NILSSON, J. PRAGLOWSKI & L. NILSSON, Atlas of airborne pollen grains and spores in Northern Europe, 321
Trematodon brevirostris, 218, 225; longicollis, 516
Tremolecia atrata, 407; jurana var. sublutescens, 318
Treubia insignis, 582
Tricharia melanothrix, 125, 139; santessonii, 88+, 89*, 90*, 505; triseptata, 110
Trichocolea tenera, 637; tomentella, 29+, 114, 310, 484
Trichosteleum cyparissoides, 243; fluviale, 243; papillosum, 243
Tristichium mirabile, 221, 630
Trichostomopsis australasiae, 227, 246, 247, 255; brevifolia, 266; crispifolia, 520; diaphanobasis, 266; fayae, 266, 544; umbrosa, 495
Trichostomum brachydontium, 227, 439, 520, 546, subsp. mutabile, 546; caespitosum, 546; campylocarpum, 255; crispulum, 496, 546; filescens, 226, 255; jamaicense, 437, 479, 486, 541; luteum, 255; novogranatense, 227; orthotrichoides, 226, 256; platyphyllum, 546; raapii, 227; schlimii, 227, 255; spirale, 546; subcirrhifolium, 227, 256; syntrichioides, 227, 256; tenuirostre, 81; triumphans, 546, var. pallidisetum, 546; viridulum, 546; wallisii, 256
Trichothecium gemmiferum, 502
Tritomaria exsectiformis, 166; polita, 118; quinquedentata, 115, 165+
TUCKER, S. New or noteworthy records of lichens from Louisiana, 125
TUCKERMAN, E. (portrait), 512*
Tuerckheimia angustifolia, 488

UDAR, R. & D. K. SINGH. Notothylas dissecta, a hornwort new to India, 625
Ulota bruchii, 557; crispa, 557; phyllantha, 498
ultrastructural localization of polysaccharides in Usnea cavernosa, 533
Umbilicaria caroliniana, 407; cinereorufescens, 407; cylindrica, 407; decussata, 487; hirsuta, 407; hyperborea, 407; krascheninnikovii, 407; lyngei, 407; muhlenbergii, 109; proboscidea, 407; rigida, 407; scholanderi, 407; torrefacta, 24, 407; vellea, 407; virginis, 407
Uruguay, contribution to the lichen flora of, 472
Usnea canariensis, 504; cavernosa, 533+, 564; dichroa var. spinulosa, 473; goniodes, 111; hirta, 24, 107; hookeri, 504; rubiginea, 139; sorediifera, 24; subfloridana, 24
usnic acid, 154

Valdiviella pauperculata, 516
validation of Tortula herzogii Zander, 631
Venezuela, new records for the moss flora of, 618
Verrucaria aspiciliae, 506; biatorinaria, 506; calciseda, 407; coccinearia, 506; devergescens, 396+; deversa, 407; eleganteria, 506; gemmifera, 502; granulosaria, 506; helveticorum, 506; insularis var. maior, 506; margacea, 386+; melaspora, 502; peripherica, 502; protearia, 506; sphinctrinella, 24; turpis, 520
Vesicularia amphibola, 244; inflectens, 79, 80; montagnei, 144; perviridis, 80; poeppigiana, 244; vesicularis, 481
Vestergrenopsis isidiata, 407
Vezdaea foliicola, 319
vicanicin, 303
VITT, D. H. *See* R. E. ANDRUS & D. H. VITT, 295

VITT, D. H. New taxa and new combinations in the Orthotrichaceae of Mexico. 1
VITT, D. H. & D. G. HORTON. Bryophytes new to Alberta, Yukon Territory and Northwest Territories, Canada, 409

Wadeana dendrographa, 315; minuta, 315
Warburgiella leptocarpa var. altepapillosa, 498
Washington, new and interesting hepatics from northern, 162
WEBER, W. A. *See* A. W. ARCHER & W. A. WEBER, 481
WEBER, W. A. (review), L. Rodenberg, Epilithische Vegetation in einem alten Weidegebiet auf Mittel-Öland, Schweden, 120
WEBER, W. A. Aloina bifrons (Musci: Pottiaceae) new to South America, 493
Webera papillosa, 256; spectabilis, 230, 256
Weissia atrocaulis, 545; bogotensis, 224, 256; canaliculata, 227; controversa, 67+, 227, 256, 266, 275, 417+, 438, 442, 486, 520, 540, 545, 609, 612; cucullata, 256; edentula, 545; fallax, 546; hampei, 225, 256; hedwigii, 541, 545; lindigiana, 256; lindigii, 224, 256; longidens, 546; microstoma, 541; newcomeri, 545; planifolia, 546; rostellata, 545, 555; rutilans, 546, var. ganderi, 546; squarrosa, 545; subcaespitosa, 256; tortilis, 545; tortula, 256; tyrrena, 546; viridula, 256, 421, var. canaliculata, 227; wimmeriana, 546
Weisiopsis anomala, 543
WESSELS, D. C., L. A. WESSELS & W. H. HOLZAPFEL. Preliminary report on lichen-feeding Coleoptera occurring on Teloschistes capensis in the Namib Desert, South West Africa, 270
West Himalayan mosses, cytological investigations in, 583
West Virginia, Tortula ammonsiana, a new species from, 469
Weymouthia orbiculata, 518
Wijkia scabrifolia, 498; subnitida, 243
Wijkiella kenyae, 494
WILLIAMS, R. S. (portrait), 512*
Wisconsin, bryophytes of southwestern, 273
WYATT, R. *See* A. STONEBURNER & R. WYATT, 491
Wyoming, additions to the mosses of, 72

Xanthoparmelia ajoensis, 156+; arida, 501; dierythra, 159; kurokawae, 157+; lecanorica, 501; lineola, 24; mexicana, 24, 157; **moctezumensis**, 154+, 156*; plittii, 159; quintaria, 108; scabrosa, 108; tucsonensis, 154
Xanthoria candelaria, 139; elegans, 24, 407; fallax, 24, 107; parietina, 106, 319, 473, 533, 536; polycarpa, 24; sorediata, 24
xanthorin, 302
Xylographa abietina, 24

Yukon Territory, bryophytes new to, 409

ZANDER, R. H. Regenerated herbarium material for biosystematic and cytological studies, 323
ZANDER, R. H. Molendoa hornschuchiana new to the Americas from the Pacific Coast of Alaska, 487
ZANDER, R. H. Lectotypification of Tortula propagulosa Sharp, 510
ZANDER, R. H. Portraits of American bryologists and lichenologists, 511, 512
ZANDER, R. H. Patterns of sporophyte maturation dates in the Pottiaceae (Bryopsida), 538
ZANDER, R. H. Validation of Tortula herzogii Zander, 631
ZANDER, R. H. & W. J. HOE. Geographic disjunction and heterophylly in Tortella fragilis var. tortelloides (=Sarconeurum tortelloides), 84
ZANDER, R. H. & R. R. IRELAND, JR. Propaguliferous Ceratodon purpureus in riparian environments, 474
ZEHR, D. R. Phenology of selected bryophytes in southern Illinois, 29
Zelometeorium patens, 235; patulum, 236; recurvifolium, 136
zinc, effect on lichens, 20+; localization in aquatic bryophytes, 594
Zygodon aureus, 233, 256; campylophyllus, 233; ceratodontoides, 233, 256; conoideus, 519; denticulatus, 233, 256; fragilis, 619; goudotii, 233, 256; linguiformis, 233, 256; moritzianus, 256; nivalis, 233; obtusifolius, 233, 256; pichinchensis, 233; reinwardtii, 233, 256, 439, var. integrifolius, 233, 256; rufescens, 233; setulosus, 233; sordidus, 233; squarrosus, 233; subdenticulatus, 233, 256; stenocarpus, 233; viridissimus, 266

Volume 82, Number 3 (pages 325–512), was issued 17 October 1979

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
DALE M. J. MUELLER

Volume 82
1979

Table of Contents

ALLEN, B. H. Additional distributional records of mosses in Pennsylvania ... 289
ALLEN, B. H. *Bryoxiphium norvegicum* in Colorado 490
ALPERT, P. Desiccation of desert mosses following a summer rainstorm 65
ANDRUS, R. E. *Sphagnum bartlettianum* in the southeastern United States .. 198
ANDRUS, R. E. & D. H. VITT. Sphagnotheca Boreali-americana: Fascicle III . 295
ARCHER, A. W. & W. A. WEBER. *Baeomyces frenchianus*, an Australian endemic and its chemotaxonomy 481
BAKER, K. K., J. A. MALACHOWSKI & G. R. HOOPER. Ultrastructural localization of polysaccharides in *Usnea cavernosa* 533
BANERJEE, R. D. & S. P. SEN. Antibiotic activity of bryophytes 141
BARCLAY-ESTRUP, P. & R. J. K. RINNE. Trace element accumulation in a feather moss and in soil near a Kraft paper mill in Ontario 599
BRASSARD, G. R. Review 121
BURTON, M. A. S. Studies on zinc localization in aquatic bryophytes 594
CHURCHILL, S. P. Mosses of the Great Plains III. Additions to Nebraska and the Black Hills of South Dakota and Wyoming 72
COLE, M., F. D. BOWERS & S. FRECKMANN. Bryophytes of the Kickapoo River Valley, southwestern Wisconsin 273
CROSBY, M. R. Recent literature on mosses—113 494
CRUM, H. & L. E. ANDERSON. *Tortula ammonsiana*, a new species from West Virginia 469
CULBERSON, C. F. & H. HERTEL. Chemical and morphological analyses of the *Lecidea lithophila-plana* group (Lecideaceae) 189
CULBERSON, C. F., T. H. NASH III & A. JOHNSON. 3-α-Hydroxybarbatic acid, a new depside in chemosyndromes of some Xanthoparmeliae with β-Orcinol depsides 154
DELGADILLO M., C. Mosses and phytogeography of the *Liquidambar* forest of Mexico 432
DELGADILLO M., C. Notes on alpine mosses of Mexico 629
DIETERT, M. F. Studies on the gametophyte nutrition of the cosmopolitan species *Funaria hygrometrica* and *Weissia controversa* 417
EGAN, R. S. Recent literature on lichens—101–103 ..... 105, 314, 500
EGUNYOMI, A. & S. O. OLARINMOYE. Mosses from Nigeria II. Collections from the Obudu cattle ranch 76
FIFE, A. J. Taxonomic observations on three species of North American *Funaria* 204
FLORSCHÜTZ-DE WAARD, J. & P. A. FLORSCHÜTZ. Estudios sobre Criptógamas Colombianas III. Lista comentada de los Musgos de Colombia 215
FRAHM, J.-P. New *Campylopus* records from Mexico, including *C. sharpii* sp. nov. 623
GLIME, J. M. & D. W. ACTON. Temperature effects on assimilation and respiration in the *Fontinalis duriaei*—Periphyton association 382
GODFREY, J. D. & W. B. SCHOFIELD. New and interesting hepatics from British Columbia, Canada, and northern Washington State, U.S.A. II. 162
GRIFFIN, D., III. New records for the moss flora of Venezuela 618
HARRIS, R. C. Four species of *Thelopsis* Nyl. (Lichenized Ascomycetes) new to North America 77

HARTHILL, M. P., D. M. LONG & B. D. MISHLER. Preliminary list of southern California mosses 260
HOE, W. J. Additional new and noteworthy records for Hawaiian mosses—5 . 79
HOLCOMBE, J. W. *Sphagnum angermanicum* in New Hampshire 616
HONG, W. S. The genus *Scapania* in western North America I. Historical background 181
HORTON, D. G. *Encalypta vittiana sp. nov.* and *E. flowersiana sp. nov.* from North America 368
HORTON, D. G. Encalyptaceae Americanae Exsiccatae: An annotated guide to Fasciculus I (Number 1–10) 450
JACOBSON, D. E. & P. V. PRIOR. Additions to the moss flora of Nebraska 622
JOURNET, A. R. P. Review 508
KITAGAWA, N. & I. KANDA. Linear spore tetrads in *Conocephalum supradecompositum* 96
KUMAR, S. S. & S. K. VERMA. Cytological investigations in some west Himalayan mosses 583
KUWAHARA, Y. Notes on the genus *Steereella* 613
LAWTON, E. *Grimmia pacifica*, a new species from western North America .. 276
LONGTON, R. E. Studies on growth, reproduction and population ecology in relation to microclimate in the bipolar moss *Polytrichum alpestre* 325
MAHANDRU, M. M. & O. L. GILBERT. Chemical studies in *Fulgensia*: Structures of two new chlorodepsidones 302
MAHANDRU, M. M. & O. L. GILBERT. Norgangaleoidin, a dichlorodepsidone from *Lecanora chlarotera* 292
MAHU, M. Familias y géneros de Musgos Chilenos 513
MANUEL, M. G. Review 120
MARSH, J. E. & T. H. NASH III. Lichens in relation to the Four Corners Power Plant in New Mexico 20
MATTERI, C. M. Notas Briológicas (Musci) IV y V 99
MILLER, N. G. Review 507
MOSER, T. J., T. H. NASH III & J. W. THOMSON. Lichens of Anaktuvuk Pass, Alaska, with emphasis on the impact of caribou grazing 393
MUELLER, D. M. J. Editorial 122
MUELLER, D. M. J. Reviews 122, 322, 509
MUELLER, D. M. J. & R. E. MAGILL. Notes on Texas hepatics 298
NASH, T. H., III & L. L. SIGAL. Gross photosynthetic response of lichens to short-term ozone fumigations 280
NORRIS, D. H. & H. ROBINSON. The systematic position of *Bryobrothera crenulata* 305
O'FLAHERTY, L. M. & G. W. FREEMAN. Eleven mosses new to Illinois 609
OSARIO, H. S. Contribution to the lichen flora of Uruguay XI. Lichens from Arazatí, San José Department 472
PAOLILLO, D. J., JR. On the lipids of the sperm masses of three mosses 93
PURSELL, R. A. On *Fissidens*. Nomenclatural and distributional notes on some American species 58
PURSELL, R. A. & E. HEGEWALD. On *Fissidens* in Perú 267
RAO, K. R., N. R. KUMAR & A. N. REDDY. Studies of photosynthesis in some liverworts 286

REDFEARN, P. L., JR. Bryophytes of the Interior Highlands of North America XVI. Species of the Ozark National Forest, Sylamore Division, and adjacent areas in Stone County, Arkansas 483
REESE, W. D. Calymperaceae (Musci) from western Amazônia: Brazil and Bolivia 559
RUNDEL, P. W., G. C. BRATT & O. L. LANGE. Habitat ecology and physiological response of *Sticta filix* and *Pseudocyphellaria delisei* from Tasmania ... 171
SANFORD, G. R. Temperature related growth patterns in *Amblystegium riparium* 525
SCHERTLER, M. M. Development of the archegonium and embryo in *Lophocolea heterophylla* 576
SELKIRK, P. M. Effect of nutritional conditions on sexual reproduction in *Riccia* 37
SÉRUSIAUX, E. Foliicolous lichens from southeastern United States 88
SEYER, S. C. New mosses from Crater Lake National Park, Oregon 82
SHERIDAN, R. P. Impact of emissions from coal-fired electricity generating facilities on N_2-fixing lichens 54
SHOWMAN, R. E. Cold injury to lichens in southern Ohio 620
SIGAL, L. L. & O. C. TAYLOR. Preliminary studies of the gross photosynthetic response of lichens to peroxyacetylnitrate fumigations 564
SPIESS, L. D. Antagonism of cytokinin induced callus in *Pylaisiella selwynii* by nucleosides and cyclic nucleotides 47
STEERE, W. C. The chromosome number of *Callicostella depressa* 103
STEERE, W. C. Some mosses from British Honduras 478
STONEBURNER, A. & R. WYATT. Three Big Thicket bryophytes new to Texas .. 491
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—83–85 ..... 112, 310, 632
TAN, B. C. Review 638
TIXIER, P. A propos de quelques mousses nouvelles pour le sud de Madagascar 598
TIXIER, P. Contribution to the knowledge of the genus *Cololejeunea* VIII. Some new species of Malagasy *Cololejeunea* 602
TRAVERSE, A. Review 321
TUCKER, S. C. New or noteworthy records of lichens from Louisiana 125
UDAR, R. & D. K. SINGH. *Notothylas dissecta*, a hornwort new to India 625
VITT, D. H. New taxa and new combinations in the Orthotrichaceae of Mexico 1
VITT, D. H. & D. G. HORTON. Bryophytes new to Alberta, Yukon Territory and Northwest Territories, Canada 409
WEBER, W. A. *Aloina bifrons* (Musci: Pottiaceae) new to South America 493
WEBER, W. A. Review 120
WESSELS, D. C. J., L. A. WESSELS & W. H. HOLZAPFEL. Preliminary report on lichen-feeding Coleoptera occurring on *Teloschistes capensis* in the Namib Desert, South West Africa 270
ZANDER, R. H. *Molendoa hornschuchiana* new to the Americas from the Pacific Coast of Alaska 487
ZANDER, R. H. Patterns of sporophyte maturation dates in the Pottiaceae (Bryopsida) 538

ZANDER, R. H. Validation of *Tortula herzogii* Zander 631
ZANDER, R. H. & W. J. HOE. Geographic disjunction and heterophylly in *Tortella fragilis* var. *tortelloides* (=*Sarconeurum tortelloides*) 84
ZANDER, R. H. & R. R. IRELAND, JR. Propaguliferous *Ceratodon purpureus* in riparian environments 474
ZEHR, D. R. Phenology of selected bryophytes in southern Illinois 29